

Appl. No. : 10/063,599
Filed : May 3, 2002

REMARKS

Applicants thank the Examiner for her review of the instant application. Claims 1-5 remain pending and are presented for further examination. For the reasons stated below, Applicants respectfully traverse the rejection of the pending claims.

Rejection Under 35 U.S.C. §101

The PTO maintains its rejection of Claims 1-5 under 35 U.S.C. § 101 as lacking a specific and substantial asserted utility or a well established utility for the reasons set forth in the previous Office Actions. The PTO asserts that there is no correlation between increased transcript levels and increased protein levels, and newly cites Anderson *et al.*, Lian *et al.*, and Fessler *et al.* as support. The PTO also relies on Hu *et al.* for support for its statement that “the literature cautions researchers from drawing conclusions based on small changes in transcript expression levels between normal and cancerous tissues.” *Final Office Action* at 6. The PTO argues that the asserted utility for the polynucleotides and polypeptides is not in currently available form, making the asserted utility not substantial. *Final Office Action* at 8

Applicants have previously set forth the legal standard for utility. It is established that the legal standard for demonstrating utility is a relatively low hurdle. An Applicant need only provide evidence such that it is **more likely than not that a person of skill in the art would be convinced, to a reasonable probability, that the asserted utility is true.** The evidence need not be direct evidence, so long as there is a reasonable correlation between the evidence and the asserted utility. The Applicant **does not need to provide evidence such that it establishes an asserted utility as a matter of statistical certainty.**

Even assuming that the PTO has met its initial burden to offer evidence that one of ordinary skill in the art would reasonably doubt the truth of the asserted utility, Applicants assert that they have met their burden of providing rebuttal evidence such that it is more likely than not those skilled in the art, to a reasonable probability, would believe that the claimed invention is useful as a diagnostic tool for cancer.

Substantial Utility

Summary of Applicants' Arguments and the PTO's Response

Applicants first offer a summary of their argument and the disputed issues involved. Applicants assert that the claimed polypeptides have utility as diagnostic tools for cancer, particularly esophageal, stomach, lung, rectal and melanoma tumors. Applicants' asserted utility rests on the following argument:

1. Applicants have provided reliable evidence that mRNA for the PRO1327 polypeptide is expressed at least two-fold higher in normal esophageal, stomach, lung, rectal and skin lung tissue compared to tumor tissue of these same types;

2. Applicants assert that it is well-established in the art that a change in the level of mRNA for a particular protein, e.g. a decrease, generally leads to a corresponding change in the level of the encoded protein, e.g. a decrease;

3. Given Applicants' evidence that the mRNA for the PRO1327 polypeptide is differentially expressed in esophageal, stomach, lung, rectal and melanoma tumor tissue compared to normal tissue of these same types, it is more likely than not that the PRO1327 polypeptide is likewise differentially expressed in these tumors, and antibodies that specifically bind the PRO1327 polypeptide are therefore useful as diagnostic tools to distinguish esophageal, stomach, lung, rectal and melanoma tumor tissue from normal tissue of these same types.

Applicants understand the PTO to be asserting that "one skilled in the art would not assume that an increase in mRNA levels results in increased protein levels without doing the empirical experimentation necessary to measure protein levels. The requirement for such empirical experimentation indicates that the asserted utility for the claimed polypeptides is not substantial; it is not in currently available form." *Final Office Action* at 5.

As detailed below, Applicants submit that the PTO has failed to demonstrate that this is one of the "rare cases" where the applicants have "asserted a utility that could only be true if it violated a scientific principle, such as the second law of thermodynamics, or a law of nature, or was wholly inconsistent with contemporary knowledge in the art." *M.P.E.P.* § 2107.02 III B. The references cited by the PTO in support of its rejections are either irrelevant, not contrary to Applicants' arguments, or actually offer support for Applicants' position. Even if the PTO has met its initial burden, Applicants have submitted enough rebuttal evidence such that it is **more**

Appl. No. : 10/063,599
Filed : May 3, 2002

likely than not that a person of skill in the art would be convinced, to a reasonable probability, that the asserted utility is true. As stated above, Applicants' evidence need not be direct evidence, so long as there is a reasonable correlation between the evidence and the asserted utility. **The standard is not absolute certainty.**

Applicants have established that the Gene Encoding the PRO1327 Polypeptide is Differentially Expressed in Certain Cancers compared to Normal Tissue

Applicants remind the PTO of the level of evidence required to support a substantial utility.

[T]he Appellant does not have to provide evidence sufficient to establish that an asserted utility is true "beyond a reasonable doubt." Nor must the Appellant provide evidence such that it establishes an asserted utility as a matter of statistical certainty. Instead, evidence will be sufficient if, considered as a whole, it leads a person of ordinary skill in the art to conclude that the asserted utility is more likely than not true. M.P.E.P. at § 2107.02, part VII (emphasis in original, citations omitted).

The Court of Appeals for the Federal Circuit has stated that the standard for satisfying the utility requirement is a low one:

The threshold of utility is not high: An invention is "useful" under section 101 if it is capable of providing some identifiable benefit. *See Brenner v. Manson*, 383 U.S. 519, 534, 86 S.Ct. 1033, 16 L.Ed.2d 69 (1966); *Brooktree Corp. v. Advanced Micro Devices, Inc.*, 977 F.2d 1555, 1571 (Fed. Cir. 1992) ("To violate § 101 the claimed device must be totally incapable of achieving a useful result"); *Fuller v. Berger*, 120 F. 274, 275 (7th Cir.1903) (test for utility is whether invention "is incapable of serving any beneficial end"). *Juicy Whip, Inc. v. Orange Bang, Inc.*, 185 F.3d 1364, 1366, 51 U.S.P.Q. 2d 1700 (Fed. Cir. 1999) (emphasis added).

The low threshold for satisfying the utility requirement is reflected in the standard set by the Federal Circuit for invalidating a patent based on a lack of utility: "[T]he fact that an invention has only limited utility and is only operable in certain applications is not grounds for finding lack of utility. Some degree of utility is sufficient for patentability. Further, the defense of non-utility cannot be sustained without proof of total incapacity." *Envirotech Corp. v. Al George, Inc.*, 730 F.2d 753, 762, 221 U.S.P.Q. 473 (Fed. Cir. 1984) (emphasis added, citations omitted).

Because the standard for satisfying the utility requirement is so low, requiring total incapacity for a finding of no utility, the M.P.E.P. cautions that:

Appl. No. : 10/063,599
Filed : May 3, 2002

Rejections under 35 U.S.C. 101 have been *rarely* sustained by federal courts. Generally speaking, in these *rare* cases, the 35 U.S.C. 101 rejection was sustained [] because the Appellant ... asserted a utility that could only be true if it violated a scientific principle, such as the second law of thermodynamics, or a law of nature, or was wholly inconsistent with contemporary knowledge in the art. M.P.E.P. § 2107.02 III B., citing *In re Gazave*, 379 F.2d 973, 978, 154 U.S.P.Q. 92, 96 (C.C.P.A. 1967) (underline emphasis in original, italic emphasis added).

In *Nelson v. Bowler*, 626 F.2d 853, 206 U.S.P.Q. 881 (C.C.P.A. 1980), the court held that crude screens for pharmacological activity which were reported as qualitative results without statistical analysis were sufficient to establish utility. The Appellants in *Nelson* relied on two tests to prove practical utility for derivatives of naturally occurring prostaglandins: an *in vivo* rat blood pressure (BP) test and an *in vitro* gerbil colon smooth muscle stimulation (GC-SMS) test. In the BP test, responses to the compounds were categorized qualitatively, as either a depressor (lowering) effect or a pressor (elevating) effect. *Nelson*, 626 F.2d at 854-55. In the GC-SMS test a section of colon was excised from a freshly-killed gerbil for suspension in a physiological solution, and a lever arm was connected to the colon in such a way that any contraction was recorded as a polygraph trace. *Id.* The Board held that Nelson had not shown adequate proof of practical utility, characterizing the tests as “rough screens, uncorrelated with actual utility.” *Id.* at 856.

On appeal the C.C.P.A. reversed, holding that the Board “erred in not recognizing that tests evidencing pharmacological activity may manifest a practical utility even though they may not establish a specific therapeutic use.” *Id.* (emphasis added). The Court stated that “practical utility” was characterized as a use of the claimed discovery in a manner which provides some immediate benefit to the public, establishing the rule that “[k]nowledge of the pharmacological activity of any compound is obviously beneficial to the public.... [W]e conclude that adequate proof of any such activity constitutes a showing of practical utility.” *Id.* (emphasis added).

The Court rejected Bowler’s argument that the BP and GC-SMS tests are inconclusive showings of pharmacological activity since confirmation by statistically significant means did not occur until after the critical date. The Court stated that “a rigorous correlation is not necessary where the test for pharmacological activity is reasonably indicative of the desired response.” *Id.* (emphasis added). The Court concluded that a “reasonable correlation” between the observed properties and the suggested use was sufficient to establish practical utility. *Id.* at 857.

The test articulated in *Nelson* is certainly met by the evidence in Example 18. Presented with the data in Example 18, one of skill in the art would find that there is a “reasonable correlation” between the observed property of differential expression in certain tumors and the suggested use as a diagnostic tool for cancer. In *Nelson* the fact that the results were qualitative, not statistically significant, and preformed *in vivo* in rats or *in vitro* on gerbil colon did not matter. The Court held that statistically significant results are not required, nor is it necessary to prove actual clinical therapeutic usefulness.

The gene expression data in the specification, Example 18, shows that the mRNA associated with the PRO1327 polypeptide was more highly expressed in normal esophageal, stomach, lung, rectal and skin tissue compared to esophageal, stomach, lung, rectal and melanoma tumor tissue. Gene expression was analyzed using standard semi-quantitative PCR amplification reactions of cDNA libraries isolated from different human tumor and normal human tissue samples. Identification of the differential expression of the PRO1327 polypeptide-encoding gene in tumor tissue compared to the corresponding normal tissue renders the molecule useful as a diagnostic tool for the determination of the presence or absence of tumor. Applicants previously submitted a first Declaration of J. Christopher Grimaldi, an expert in the field of cancer biology. This declaration explains the importance of the data in Example 18, and how differential gene and protein expression studies are used to differentiate between normal and tumor tissue (see Declaration, paragraph 7).

In paragraph 5 of his declaration, Mr. Grimaldi states that the gene expression studies reported in Example 18 of the instant application were made from pooled samples of normal and of tumor tissues. Mr. Grimaldi explains that:

The DNA libraries used in the gene expression studies were made from pooled samples of normal and of tumor tissues. *Data from pooled samples is more likely to be accurate than data obtained from a sample from a single individual.* That is, the detection of variations in gene expression is likely to represent a more generally relevant condition when pooled samples from normal tissues are compared with pooled samples from tumors in the same tissue type. (Paragraph 5) (emphasis added).

Thus, the use of pooled samples increases the accuracy of the experiment. As Dr. Grimaldi explained, the detection of variations in gene expression is likely to represent a more generally relevant condition when pooled samples from normal tissues are compared with pooled

Appl. No. : **10/063,599**
Filed : **May 3, 2002**

samples from tumors in the same tissue type. Clinical diagnostics are geared towards generally relevant conditions that are present in a populous.

With respect to the PTO's concerns regarding the methodology used to compare mRNA levels in normal tissue to that in cancerous tissue in Example 18, Applicants maintain that this methodology is reliable. In paragraphs 6 and 7, Mr. Grimaldi explains that the semi-quantitative analysis employed to generate the data of Example 18 is sufficient to determine if a gene is over- or under-expressed in tumor cells compared to corresponding normal tissue. He states that any visually detectable difference seen between two samples is indicative of at least a two-fold difference in cDNA between the tumor tissue and the counterpart normal tissue. Thus, the results of Example 18 reflect at least a two-fold difference between normal and tumor samples. He also states that the results of the gene expression studies indicate that the genes of interest "can be used to differentiate tumor from normal," thus establishing their reliability. He explains that, "The precise levels of gene expression are irrelevant; what matters is that there is a relative difference in expression between normal tissue and tumor tissue." (Paragraph 7). Thus, since it is the relative level of expression between normal tissue and suspected cancerous tissue that is important, the precise level of expression in normal tissue is irrelevant. Likewise, there is no need for quantitative data to compare the level of expression in normal and tumor tissue. As Mr. Grimaldi states, "If a difference is detected, this indicates that the gene and its corresponding polypeptide and antibodies against the polypeptide are useful for diagnostic purposes, to screen samples to differentiate between normal and tumor."

With respect to the Examiner's assertions that Mr. Grimaldi's Declaration is unpersuasive because, as an inventor and employee of the assignee, Mr. Grimaldi has an interest in the outcome of the case, Applicants note that an affidavit cannot be disregarded solely because it is signed by the applicant. (See M.P.E.P. §716.01(c)). Furthermore, Applicants maintain that Mr. Grimaldi's first Declaration objectively sets forth the methodology employed in the experiments described in Example 18 and the conclusions derived therefrom, and Mr. Grimaldi's second Declaration objectively sets forth the understanding of those skilled in the art regarding the correlation between differential mRNA expression and differential polypeptide expression. In addition, declarations relating to issues of fact should not be summarily dismissed without an

Appl. No. : 10/063,599
Filed : May 3, 2002

adequate explanation of how the declaration fails to rebut the Examiner's position. *See in re Alton* 76 F.3d 1168 (Fed. Cir. 1996).

Applicants turn next to the PTO's arguments based on *Hu et al.* Applicants have discussed this reference at length in its previous responses. In addition to the persuasive reasons articulated in Applicants' arguments of record, the PTO's reliance on *Hu* is also misplaced because Applicants are not relying on microarray data as discussed in *Hu*:

In any microarray experiment, thousands of genes may demonstrate statistically significant expression changes, but only a fraction of these may be relevant to the study. *Hu* at 405, left column, first paragraph (emphasis added).

Instead, Applicants are relying on a more accurate and reliable method of assessing changes in mRNA level, namely quantitative PCR analysis. In a recent study by Kuo *et al.*, (Proteomics 5(4):894-906 (2005)), the authors used microarray analysis combined with proteomic analysis using two-dimensional gel electrophoresis to examine changes in gene expression in leukemia cell lines. The authors report that "[c]omparison of microarray and proteomic expression profiles showed poor correlation. Use of more reliable and sensitive analyses, such as reverse transcriptase polymerase chain reaction [RT-PCR], Western blotting and functional assays, on several genes and proteins, nonetheless, confirmed that there is indeed good correlation between mRNA and protein expression." Kuo *et al.* at Abstract (emphasis added) (attached as Exhibit 1). Thus, even if accurate, *Hu*'s statements regarding microarray studies are not relevant to the instant application which does not rely on microarray data.

In sum, the data in Example 18 are sufficient to establish a practical utility for the claimed invention. Applicants are asserting that the PRO1327 gene, polypeptide and antibodies have utility as diagnostic tools for cancer, particularly esophageal, stomach, lung, rectal and melanoma cancer. Applicants are not asserting that the PRO1327 gene, polypeptide and antibodies necessarily provide a definitive diagnosis of cancer, but rather that they are useful, alone or in combination with other diagnostic tools, to assist in the diagnosis of cancer. Statistically significant results are not required, nor is it necessary to prove actual clinical therapeutic usefulness.

Applicants have established that the Accepted Understanding in the Art is that there is a Positive Correlation between Changes in mRNA Levels and Changes in the Level of Expression of the Encoded Protein

Applicants next turn to the second portion of their argument in support of their asserted utility – that it is well-established in the art that a change in the level of mRNA for a particular protein, generally leads to a corresponding change in the level of the encoded protein; given Applicants' evidence of differential expression of the mRNA for the PRO1327 polypeptide in esophageal, stomach, lung, rectal and melanoma tumors, it is more likely than not that the PRO1327 polypeptide is likewise differentially expressed in these tumors; and antibodies to proteins differentially expressed in certain tumors have utility as diagnostic tools.

The PTO's cited references are not contrary to Applicants' asserted utility

The PTO relies on Haynes *et al.* and Gygi *et al.*, as well as newly cited references by Anderson and Seilhamer, Lian *et al.*, and Fessler *et al.*

Haynes *et al.*, Gygi *et al.*, and Anderson *et al.*

Applicants have previously discussed at length why the Haynes *et al.* and Gygi *et al.* references are not relevant to the issue of whether changes in mRNA level for a particular gene lead to changes in protein level. Briefly stated, references such as Haynes, Gygi, and Anderson, which discuss the correlation between static levels of mRNA and static levels of protein across different genes, are not relevant – Applicants rely only on the assertion that changes in mRNA level generally lead to corresponding changes in the encoded protein level. Applicants incorporate by reference the previous arguments made regarding these issues, and will not repeat them here.

However, in an attempt to illustrate why references which relate to static global levels of mRNA and protein across different genes are not relevant to Applicants' asserted utility, Applicants offer the following illustration and analogy with the understanding that like all illustrations and analogies, they are not perfect and therefore do not represent any admissions or binding statements regarding Applicants' disclosure or invention.

Haynes, Gygi, and Anderson all discuss whether there is a correlation between the static level of mRNAs and proteins globally, *i.e.* across different genes. This is equivalent to conducting a hypothetical Experiment 1, where a particular cell type has 100 copies of mRNA

for gene X, 200 copies of mRNA for gene Y, and 400 copies of mRNA for gene Z. If there is a global correlation between static mRNA levels and protein levels across genes, the ratio of the amount of proteins X:Y:Z would be approximately 1:2:4. This is essentially what the cited references examined.

In contrast, Applicants are relying on a correlation between changes in mRNA level for a particular gene leading to a corresponding change in the level of the encoded protein. For example, in hypothetical Experiment 2, if gene X has 100 copies of mRNA per cell in condition A (*e.g.* normal), and 200 copies of mRNA for gene X in condition B (*e.g.* tumor), the ratio of the amount of protein X in condition A:B would be approximately 1:2, such that there is a correlation between the change in the level of mRNA and protein for a particular gene.

The PTO argues that because there is no correlation between static levels of mRNA and protein across genes, as illustrated by Experiment 1, one of skill in the art would not expect an increase or decrease in the amount of mRNA for a particular gene to result in a corresponding change in the amount of the encoded protein, as illustrated in Experiment 2. This is simply wrong.

For example, Haynes reports that the amount of protein produced by similar levels of mRNA varied by as much as fifty-fold, and that similar amounts of protein were sustained by amounts of mRNA that varied by as much as forty-fold. *Haynes* at 1863, first full paragraph. Based on these results, Haynes concludes that “protein levels cannot be accurately predicted from the level of the corresponding mRNA transcript.” *Id.*

This is analogous to a finding that on one gallon of gas, a hybrid car can travel 70 miles but a large truck can only travel 5 miles, or that to travel 70 miles, a hybrid car requires 1 gallon of gas, but a large truck requires 14 gallons. That is to say, there are many things which affect the fuel efficiency of a car. Based on these observations, one could conclude that given the lack of correlation between the amount of gas in an automobile and the distance it travels, one cannot predict how far an automobile will travel based on the amount of gas in the tank.

Even if true, Haynes’ data and conclusions are irrelevant to Applicants’ assertion, which is that increasing or decreasing the amount of mRNA for a particular gene will result in a corresponding increase or decrease in the amount of the encoded protein. This is analogous to increasing or decreasing the amount of gas in an automobile – it will travel farther if you add

Appl. No. : 10/063,599
Filed : May 3, 2002

more gas, and not as far with less. The fact that there are many things which affect fuel efficiency and therefore you cannot predict how far an automobile will travel without knowing if it is a hybrid or a large truck is irrelevant – both a hybrid and a truck travel farther on more gas, and not as far on less.

Applicants emphasize, and the PTO will recognize, that these are simplified illustrations to demonstrate the difference between the two issues being examined. However, these illustrations make clear that even if there is no correlation in the first experiment looking at static levels of mRNA and protein across different genes, there can still be a correlation between changes in mRNA and protein for a particular gene as examined in the second experiment. As these illustrations make clear, the PTO's evidence simply is not relevant to answering the question of whether it is **more likely than not** that a person of skill in the art would be convinced, **to a reasonable probability**, that the asserted utility is true.

Lian et al.

The PTO cites Lian *et al.* as showing a lack of correlation between mRNA expression and protein expression in mouse cells. *Final Office Action* at 4.

In Lian, the authors looked at the mRNA and protein levels of genes in a derived promyelocytic mouse cell-line during differentiation of the cells from a promyelocytic stage of development to mature neutrophils following treatment with retinoic acid. Lian at Abstract. The level of mRNA expression was measured using 3'-end differential display (DD) and oligonucleotide chip array hybridization, and protein levels were qualitatively assessed following 2-dimensional gel electrophoresis. *Id.* at Abstract, Table 6.

Lian *et al.* used DD and array hybridization to examine the expression of genes 0, 24, 48 and 72 hours after treatment with retinoic acid. *Id.* at 515, col. 1, ¶ 2. Using this information, the authors constructed a database of mRNA level changes during differentiation of the cell line. *Id.* at 518, col. 2, ¶ 2. Lian *et al.* also examined protein expression at 0 and 72 hours after retinoic acid treatment. Lian reports that they were able to identify 28 proteins which they considered differentially expressed. *Id.* at 521, Fig. 5. Of those 28, only 18 had corresponding gene expression information in the database, and only 13 had measurable levels of mRNA expression. *Id.* at 521, Table 6. The authors then compared the qualitative protein level from the 2-D electrophoresis gel to the corresponding mRNA level, and reported that only 4 genes of the 18

Appl. No. : 10/063,599
Filed : May 3, 2002

present in the database had expression levels which were consistent with protein levels. *Id.* at 512, col. 1. The authors note that “[n]one of these was on the list of genes that were differentially expressed significantly (5-fold or greater change by array or 2-fold or greater change by DD).” *Id.* at 512, bridge paragraph (emphasis added). Based on these data, the authors conclude “[f]or protein levels based on estimated intensity of Coomassie dye staining in 2DE, there was poor correlation between changes in mRNA levels and estimated protein levels.” *Id.* at 522, col. 2, ¶ 2.

These results are not contrary to Applicants’ assertion. Applicants again emphasize that Applicants are asserting that a measurable change in mRNA level generally leads to a corresponding change in the level of protein expression, not that changes in protein level can be used to predict changes in mRNA level. Based on the authors’ criteria, mRNA levels were significantly changed if they were at least 5-fold different when measured using a microchip array, or 2-fold different when using the more sensitive 3’-end differential display (DD). Of the 28 proteins listed in Table 6, only one has an mRNA level measured by microarray which is differentially expressed according to the authors (spot 7: melanoma X-actin, which mRNA changed from 2539 to 341.3, and protein changed from 1 to 3). None of the other mRNAs listed in Table 6 show a significant change in expression level when using the criteria established by the authors for the less sensitive microarray technique.

There is also one gene in Table 6 whose expression was measured by the more sensitive technique of DD, and its level increased from a qualitative value of 0 to 2, a more than 2-fold increase (spot 2: actin, gamma, cytoplasmic). This increase in mRNA was accompanied by a corresponding increase in protein level, from 3 to 6.

Therefore, although the authors characterize the mRNA and protein levels as having a “poor correlation,” this does not reflect a lack of a correlation between a change in mRNA level and a corresponding change in protein level. Only two genes meet the authors’ criteria for differentially expressed mRNA level, and of those, one apparently shows a corresponding change in protein level and one does not. *Id.* at 521, Table 6. Thus, there is little basis for the authors’ conclusion relied on by the PTO that “it may be difficult to extrapolate directly from individual mRNA changes to corresponding ones in protein levels.” *Final Office Action* at 4 (emphasis added).

Fessler et al.

The PTO also cites a publication by Fessler *et al.* Fessler is not contrary to Applicants' asserted utility, and actually supports Applicants' assertion that a change in the level of mRNA for a particular protein generally leads to a corresponding change in the level of the encoded protein. As noted above, Applicants make no assertions regarding changes in protein levels when mRNA levels are unchanged, nor does evidence of changes in protein levels when mRNA levels are unchanged have any relevance to Applicants' asserted utility.

Fessler *et al.* studied changes in neutrophil (PMN) gene transcription and protein expression following lipopolysaccharide (LPS) exposure. Fessler lists in Table VIII a comparison of the change in the level of mRNA for 13 up-regulated proteins and 5 down-regulated proteins. Of the 13 up-regulated proteins, a change in mRNA levels is reported for only 3 such proteins. For these 3, mRNA levels are increased in 2 and decreased in the third. Of the 5 down-regulated proteins, a change in mRNA is reported for 3 such proteins. In all 3, mRNA levels also are decreased. Thus, in 5 of the 6 cases for which a change in mRNA levels are reported, the change in the level of mRNA corresponds to the change in the level of the protein. This is consistent with Applicants' assertion that a change in the level of mRNA for a particular protein generally leads to a corresponding change in the level of the encoded protein.

Regarding the remainder of the proteins listed in Table VIII, in 6 instances, protein levels changed while mRNA levels were unchanged. This evidence has no relevance to Applicants' assertion that changes in mRNA levels lead to corresponding changes in protein levels, since Applicants are not asserting that changes in mRNA levels are the only cause of changes in protein levels. In the final 6 instances listed in Table VIII, protein levels changed while mRNA was noted as "absent." This evidence also has no relevance to Applicants' assertion that changes in mRNA levels causes corresponding changes in protein levels. By virtue of being "absent," it is not possible to tell whether mRNA levels were increased, decreased or remained unchanged in PMN upon contact with LPS. Nothing in these results by Fessler suggests that a change in the level of mRNA for a particular protein does not generally lead to a corresponding change in the level of the encoded protein. Accordingly, these results are not contrary to Applicants' assertions.

Appl. No. : 10/063,599
Filed : May 3, 2002

The PTO points to Fessler's statement regarding Table VIII that found a "poor concordance between mRNA transcript and protein expression changes." *Final Office Action* at 4. As is clear from the above discussion, this statement does not relate to a lack of correlation between a change in mRNA levels leading to a change in protein levels, because in 5 of 6 such instances, changes in mRNA and protein levels correlated well. Instead, this statement relates to observations in which protein levels changed when mRNA was either unchanged or "absent." As such, this statement is an observation that in addition to transcriptional activity, LPS also has post-transcriptional and possibly post-translational activity that affect protein levels, an observation which is not contrary to Applicants' assertions. Accordingly, Fessler's results are consistent with Applicants' assertion that a change in mRNA level of for a particular protein generally leads to a corresponding change in the level of the encoded protein, since 5 of 6 genes demonstrated such a correlation.

Taken as a whole, the references cited by the PTO do not support the PTO's rejection of Applicants' assertion that more often than not, there is a correlation between changes in mRNA level and changes in the level of the corresponding protein. If anything, the cited references support Applicants' position.

Applicants' previously submitted supporting declarations and references

In support of the assertion that changes in mRNA are positively correlated to changes in protein levels, Applicants previously submitted a copy of a second Declaration by J. Christopher Grimaldi, a copy of the declaration of Paul Polakis, Ph.D., excerpts from Molecular Biology of the Cell, a leading textbook in the field (Bruce Alberts, *et al.*, Molecular Biology of the Cell (3rd ed. 1994), and (4th ed. 2002), excerpts from the textbook, Genes VI, (Benjamin Lewin, Genes VI (1997)), a reference by Zhigang *et al.*, World Journal of Surgical Oncology 2:13, 2004, and a reference by Meric *et al.*, Molecular Cancer Therapeutics, vol. 1, 971-979 (2002). The details of these teachings, and how they support Applicants' asserted utility, are of record and will not be repeated here.

Applicants' additional supporting references

In addition to the supporting references previously submitted by Applicants, Applicants submit the following references to further support the assertion that changes in mRNA levels generally lead to corresponding changes in the level of the encoded polypeptide.

In a comprehensive study by Orntoft *et al.* (Mol. Cell. Proteomics. 2002; 1(1):37-45) (previously submitted with IDS, attached hereto as Exhibit 2), the authors examined gene amplification, mRNA expression level, and protein expression in pairs of non-invasive and invasive human bladder tumors. *Id.* at Abstract. The authors examined 40 well resolved abundant known proteins, and found that “[i]n general there was a highly significant correlation ($p < 0.005$) between mRNA and protein alterations. Only one gene showed disagreement between transcript alteration and protein alteration.” *Id.* at 42, col. 2. The alternations in mRNA and protein included both increases and decreases. *Id.* at 43, Table II. Clearly, a correlation in 39 of 40 genes examined supports Applicants’ assertion that changes in mRNA level generally lead to corresponding changes in protein level.

In a study by Wang *et al.* (Urol. Res. 2000; 28(5):308-15) (abstract attached as Exhibit 3) the authors report that down-regulation of E-cadherin protein has been shown in various human tumors. *Id.* at Abstract. In the reported study, the authors examined the expression of cadherins and associated catenins at the mRNA level in paired tumor and nonneoplastic primary prostate cultures. They report that “[s]ix of seven cases of neoplastic cultures showed moderately-to-markedly decreased levels of E-cadherin and P-cadherin mRNA. Similar losses of alpha-catenin and beta-catenin mRNA were also observed.” *Id.* As Applicants’ assertion would predict, the authors state that the mRNA measures showed “good correlation” with the results from protein measures. The authors conclude by stating that “this paper presents a coordinated down-regulation in the expression of E-cadherin and associated catenins at the mRNA and protein level in most of the cases studied.” *Id.*

In a more recent study by Munaut *et al.* (Int. J. Cancer. 2003; 106(6):848-55) (abstract attached as Exhibit 4) the authors report that vascular endothelial growth factor (VEGF) is expressed in 64-95% of glioblastomas (GBMs), and that VEGF receptors (VEGFR-1, its soluble form sVEGFR-1, VEGFR-2 and neuropilin-1) are expressed predominantly by endothelial cells. *Id.* at Abstract. The authors explain that infiltrating tumor cells and newly-formed capillaries progress through the extracellular matrix by local proteolysis involving matrix metalloproteinases (MMPs). In the present study, the authors “used quantitative RT-PCR, Western blot, gelatin zymography and immunohistochemistry to study the expression of VEGF, VEGFR-1, VEGFR-2, sVEGFR-1, neuropilin-1, MT1-MMP, MMP-2, MMP-9 and TIMP-2 in

20 human GBMs and 5 normal brains. The expression of these MMPs was markedly increased in most GBMs with excellent correlation between mRNA and protein levels.” *Id.* Thus, the results support Applicants’ assertion that changes in mRNA level lead to corresponding changes in protein level.

In another recent study, Hui *et al.* (Leuk. Lymphoma. 2003; 44(8):1385-94 (abstract attached as Exhibit 5) used real-time quantitative PCR and immunohistochemistry to evaluate cyclin D1 mRNA and protein expression levels in mantle cell lymphoma (MCL). *Id.* at Abstract. The authors report that seven of nine cases of possible MCL showed overexpression of cyclin D1 mRNA, while two cases showed no cyclin D1 mRNA increase. *Id.* Similarly, “[s]ix of the seven cyclin D1 mRNA overexpressing cases showed increased cyclin D1 protein on tissue array immunohistochemistry; one was technically suboptimal.” *Id.* The authors conclude that the study “demonstrates good correlation and comparability between measure of cyclin D1 mRNA ... and cyclin D1 protein.” *Id.* Thus, this reference supports Applicants’ assertion.

In a recent study by Khal *et al.* (Int. J. Biochem. Cell Biol. 2005; 37(10):2196-206) (abstract attached as Exhibit 6) the authors report that atrophy of skeletal muscle is common in patients with cancer and results in increased morbidity and mortality. *Id.* at Abstract. To further understand the underlying mechanism, the authors studied the expression of the ubiquitin-proteasome pathway in cancer patient muscle using a competitive RT-PCR to measure expression of mRNA for proteasome subunits C2 and C5, while protein expression was determined by western blotting. “Overall, both C2 and C5 gene expression was increased by about three-fold in skeletal muscle of cachectic cancer patients (average weight loss 14.5+/-2.5%), compared with that in patients without weight loss, with or without cancer. ... There was a good correlation between expression of proteasome 20Salpha subunits, detected by western blotting, and C2 and C5 mRNA, showing that increased gene expression resulted in increased protein synthesis.” These findings support Applicants’ assertion that changes in mRNA level lead to changes in protein level.

Maruyama *et al.* (Am. J. Patho. 1999; 155(3):815-22) (abstract attached as Exhibit 7) investigated the expression of three Id proteins (Id-1, Id-2 and Id-3) in normal pancreas, in pancreatic cancer and in chronic pancreatitis (CP). The authors report that pancreatic cancer cell lines frequently coexpressed all three Ids, “exhibiting good correlation between Id mRNA and

Appl. No. : 10/063,599
Filed : May 3, 2002

protein levels.” *Id.* at Abstract. In addition, the authors teach that all three Id mRNA levels were expressed at high levels in pancreatic cancer samples compared to normal or CP samples. At the protein level, Id-1 and Id-2 staining was faint in normal tissue, while Id-3 ranged from weak to strong. In contrast, in the cancer tissues “many of the cancer cells exhibited abundant Id-1, Id-2, and Id-3 immunoreactivity,” and Id-1 and Id-2 protein was increased significantly in the cancer cells by comparison to the respective controls, mirroring the overexpression at the mRNA level. Thus, the authors report that in both cell lines and tissue samples, increased mRNA levels leads to an increase in protein overexpression, supporting Applicants’ assertion.

Support for Applicants’ assertion is also found in an article by Caberlotto *et al.* (Neurosci. Lett. 1999; 256(3):191-4) (abstract attached as Exhibit 8). In a previous study, the authors investigated alterations of neuropeptide Y (NPY) mRNA expression in the Flinders Sensitive Line rats (FSL), an animal model of depression. *Id.* at Abstract. The authors reported that in the current study, that NPY-like immunoreactivity (NPY-LI) was decreased in the hippocampal CA region, and increased in the arcuate nucleus, and that fluoxetine treatment elevated NPY-LI in the arcuate and anterior cingulate cortex. The authors state that “[t]he results demonstrate a good correlation between NPY peptide and mRNA expression.” Thus, increases and decreases in mRNA levels were reflected in corresponding changes in protein level.

Misrachi and Shemesh (Biol. Reprod. 1999; 61(3):776-84) (abstract attached as Exhibit 9) investigated their hypothesis that FSH regulates the bovine cervical prostaglandin E(2) (PGE(2)) synthesis that is known to be associated with cervical relaxation and opening at the time of estrus. *Id.* at Abstract. Cervical tissue from pre-estrous/estrous, luteal, and postovulatory cows were examined for the presence of bovine (b) FSH receptor (R) and its corresponding mRNA. The authors report that bFSHR mRNA in the cervix was maximal during pre-estrus/estrus, and that the level of FSHR protein was significantly higher in pre-estrous/estrous cervix than in other cervical tissues. *Id.* The authors state that “[t]here was a good correlation between the 75-kDa protein expression and its corresponding transcript of 2.55 kb throughout the estrous cycle as described by Northern blot analysis as well as RT-PCR.” *Id.* Thus, changes in the level of mRNA for bFSHR led to corresponding changes in FSHR protein levels, a result which supports Applicants’ assertion.

Appl. No. : 10/063,599
Filed : May 3, 2002

In a study by Stein *et al.* (J. Urol. 2000; 164(3 Pt 2):1026-30) (abstract attached as Exhibit 10), the authors studied the role of the regulation of calcium ion homeostasis in smooth muscle contractility. *Id.* at Abstract. The authors investigated the correlation between sarcoplasmic endoplasmic reticulum, calcium, magnesium, adenosine triphosphatase (SERCA) protein and gene expression, and the contractile properties in the same bladder. Partial bladder outlet obstructions were created in adult New Zealand white rabbits, which were divided into control, sham operated and obstructed groups. Stein *et al.* report that “[t]he relative intensities of signals for the Western [protein] and Northern [mRNA] blots demonstrated a strong correlation between protein and gene expression. ... The loss of SERCA protein expression is mediated by down-regulation in gene expression in the same bladder.” *Id.* This report supports Applicants’ assertion that changes in mRNA level, e.g. a decrease, lead to a corresponding change in the level of the encoded protein, e.g. a decrease.

In an article by Gou and Xie (Zhonghua Jie He He Hu Xi Za Zhi. 2002; 25(6):337-40) (abstract attached as Exhibit 11) the authors investigated the expression of macrophage migration inhibitory factor (MIF) in human acute respiratory distress syndrome(ARDS) by examining the expression of MIF mRNA and protein in lung tissue in ARDS and normal persons. *Id.* at Abstract. The authors report “undetectable or weak MIF mRNA and protein expression in normal lungs. In contrast, there was marked upregulation of MIF mRNA and protein expression in the ARDS lungs.” *Id.* This is consistent with Applicants’ assertion that a change in mRNA for a particular gene, e.g. an increase, generally leads to a corresponding change in the level of protein expression, e.g. an increase.

These studies are representative of numerous published studies which support Applicants’ assertion that changes in mRNA level generally lead to corresponding changes in the level of the expressed protein. Applicants submit herewith an addition 70 references (abstracts attached as Exhibit 12) which support Applicants’ assertion.

In addition to these supporting references, Applicants also submit herewith additional references which offer indirect support of Applicants’ asserted utility. As discussed above, Applicants have challenged the relevance of references such as Haynes *et al.*, and Gygi *et al.*, which do not attempt to examine the correlation between a change in mRNA level and a change in the level of the corresponding protein level. Because the PTO continues to rely on these

references, Applicants are submitting references which report results that are contrary to the PTO's cited references and offer indirect support for Applicants' asserted utility.

For example, in an article by Futcher *et al.* (Mol. Cell Biol. 1999; 19(11):7357-68) (abstract attached as Exhibit 13) the authors conducted a study of mRNA and protein expression in yeast which was nearly identical to the one conducted by Gygi *et al.* Contrary to the results of the earlier study by Gygi, Futcher *et al.* report "a good correlation between protein abundance, mRNA abundance, and codon bias." *Id.* at Abstract.

In a study which is more closely related to Applicants' asserted utility, Godbout *et al.* (J. Biol. Chem. 1998; 273(33):21161-8) (abstract attached as Exhibit 14) studied the DEAD box gene, DDX1, in retinoblastoma and neuroblastoma tumor cell lines. The authors report that "there is a good correlation with DDX1 gene copy number, DDX1 transcript levels, and DDX1 protein levels in all cell lines studied." *Id.* Thus, in these cancer cell lines, DDX1 mRNA and protein levels are correlated.

Similarly, in an article by Papotti *et al.* (Virchows Arch. 2002; 440(5):461-75) (abstract attached as Exhibit 15) the authors examined the expression of three somatostatin receptors (SSTR) at the mRNA and protein level in forty-six tumors. *Id.* at Abstract. The authors report a "good correlation between RT-PCR [mRNA level] and IHC [protein level] data on SSTR types 2, 3, and 5." *Id.*

Van der Wilt *et al.* (Eur. J. Cancer. 2003; 39(5):691-7) (abstract attached as Exhibit 16) studied deoxycytidine kinase (dCK) in seven cell lines, sixteen acute myeloid leukemia samples, ten human liver samples, and eleven human liver metastases of colorectal cancer origin. *Id.* at Abstract. The authors report that "enzyme activity and protein expression levels of dCK in cell lines were closely related to the mRNA expression levels" and that there was a "good correlation between the different dCK measurements in malignant cells and tumors." *Id.*

Grenback *et al.* (Regul. Pept. 2004; 117(2):127-39) (abstract attached as Exhibit 17) studied the level of galanin in human pituitary adenomas using a specific radioimmunoassay. *Id.* at Abstract. The authors report that "[i]n the tumors analyzed with in situ hybridization there was a good correlation between galanin peptide levels and galanin mRNA expression." *Id.*

Similarly, Shen *et al.* (Blood. 2004; 104(9):2936-9) (abstract attached as Exhibit 18) examined the level of B-cell lymphoma 2 (BCL2) protein expression in germinal center (GC)

Appl. No. : **10/063,599**
Filed : **May 3, 2002**

B-cells and diffuse large B-cell lymphoma (DLBCL). *Id.* at Abstract. The authors report that “GC cells had low expression commensurate with the low protein expression level” and that in DLBCL the level of BCL2 mRNA and protein expression showed “in general, a good correlation.” *Id.*

Likewise, in an article by Fu *et al.* (Blood 2005; 106(13):4315-21) (abstract attached as Exhibit 19) the authors report that six mantle cell lymphomas studied “expressed either cyclin D2 (2 cases) or cyclin D3 (4 cases).” *Id.* at Abstract. “There was a good correlation between cyclin D protein expression and the corresponding mRNA expression levels by gene expression analysis.” *Id.*

These examples are only a few of the many references Applicants could cite in rebuttal to the PTO’s arguments. Applicants submit herewith 26 additional references (abstracts attached as Exhibit 20) which also support Applicants’ assertion in that they report a correlation between the level of mRNA and corresponding protein, contrary to the assertion of the PTO that mRNA and protein levels are not correlated.

In summary, Applicants submit herewith a total of 113 references in addition to the declarations and references already of record which support Applicants’ asserted utility, either directly or indirectly. These references support the assertion that in general, a change in mRNA expression level for a particular gene leads to a corresponding change in the level of expression of the encoded protein. As Applicants have previously acknowledged, the correlation between changes in mRNA level and protein level is not exact, and there are exceptions (*see, e.g.*, abstracts attached as Exhibit 21). However, Applicants remind the PTO that the asserted utility does not have to be established to a statistical certainty, or beyond a reasonable doubt. *See M.P.E.P.* at § 2107.02, part VII (2004). Therefore, the fact that there are exceptions to the correlation between changes in mRNA and changes in protein does not provide a proper basis for rejecting Applicants’ asserted utility. Applicants submit that considering the evidence as a whole, with the overwhelming majority of the evidence supporting Applicants’ asserted utility, a person of skill in the art would conclude that Applicants’ asserted utility is “more likely than not true.” *Id.*

Applicants also previously submitted the Polakis Declaration in support of their position that in general, changes in mRNA levels correlate with changes in protein levels. Applicants

Appl. No. : 10/063,599
Filed : May 3, 2002

submit herewith as Exhibit 22 a second Declaration by Dr. Polakis (Polakis II) that presents evidentiary data in Exhibit B. Exhibit B of the Declaration identifies 28 gene transcripts out of 31 gene transcripts (i.e., greater than 90%) that showed good correlation between tumor mRNA and tumor protein levels. As Dr. Polakis' Declaration (Polakis II) says "[a]s such, in the cases where we have been able to quantitatively measure both (i) mRNA and (ii) protein levels in both (i) tumor tissue and (ii) normal tissue, we have observed that in the vast majority of cases, there is a very strong correlation between increases in mRNA expression and increases in the level of protein encoded by that mRNA." Accordingly, Dr. Polakis has provided the facts to enable the Examiner to draw independent conclusions.

The case law has clearly established that in considering affidavit evidence, the PTO must consider all of the evidence of record anew. *In re Rinehart*, 531 F.2d 1084, 189 USPQ 143 (C.C.P.A. 1976) and *In re Piasecki*, 745 F.2d 1015, 226 USPQ 881 (Fed. Cir. 1985). "After evidence or argument is submitted by the applicant in response, patentability is determined on the totality of the record, by a preponderance of the evidence with due consideration to persuasiveness of argument." *In re Alton*, 37 U.S.P.Q.2d 1578, 1584 (Fed. Cir. 1996)(quoting *In re Oetiker*, 977 F.2d 1443, 1445, 24 U.S.P.Q.2d 1443, 1444 (Fed. Cir. 1992)). Furthermore, the Federal Court of Appeals held in *In re Alton*, "We are aware of no reason why opinion evidence relating to a fact issue should not be considered by an examiner." *Id.* at 1583. Applicants also respectfully draw the PTO's attention to the Utility Examination Guidelines which state, "Office personnel must accept an opinion from a qualified expert that is based upon relevant facts whose accuracy is not being questioned; it is improper to disregard the opinion solely because of a disagreement over the significance or meaning of the facts offered." Part IIB, 66 Fed. Reg. 1098 (2001).

The PTO has asserted that the state of the art is such that polypeptide levels cannot be accurately predicted from mRNA levels. Applicants have addressed each of the PTO's supporting references and shown that they are either irrelevant, or taken as a whole, actually support Applicants' assertion that a change in mRNA level leads to a corresponding change in the level of the encoded protein. In addition, Applicants have submitted expert declarations, textbook excerpts, and over 115 scientific publications which support Applicants' arguments.

Appl. No. : 10/063,599
Filed : May 3, 2002

In conclusion, Applicants submit that they have offered sufficient evidence to establish that it is more likely than not that one of skill in the art would believe that because the PRO1327 mRNA is differentially expressed in certain tumor tissue, the PRO1327 polypeptide will likewise be differentially expressed in these tumors. This differential expression of the PRO1327 polypeptide makes the claimed antibodies useful as diagnostic tools for cancer, particularly esophageal, stomach, lung, rectal and melanoma tumor.

The Arguments made by the PTO are not Sufficient to satisfy the PTO's Initial Burden of Offering Evidence "that one of ordinary skill in the art would reasonably doubt the asserted utility"

As stated above, an Applicant's assertion of utility creates a presumption of utility that will be sufficient to satisfy the utility requirement of 35 U.S.C. § 101, "unless there is a reason for one skilled in the art to question the objective truth of the statement of utility or its scope." *In re Langer*, 503 F.2d 1380, 1391, 183 USPQ 288, 297 (CCPA 1974). The evidentiary standard to be used throughout *ex parte* examination in setting forth a rejection is a preponderance of the evidence, or "more likely than not" standard. *In re Oetiker*, 977 F.2d 1443, 1445, 24 USPQ2d 1443, 1444 (Fed. Cir. 1992). This is stated explicitly in the M.P.E.P.:

[T]he applicant does not have to provide evidence sufficient to establish that an asserted utility is true "beyond a reasonable doubt." **Nor must the applicant provide evidence such that it establishes an asserted utility as a matter of statistical certainty.** Instead, evidence will be sufficient if, considered as a whole, it leads a person of ordinary skill in the art to conclude that the asserted utility is more likely than not true. M.P.E.P. at § 2107.02, part VII (2004) (underline emphasis in original, bold emphasis added, internal citations omitted).

The PTO has the initial burden to offer evidence "that one of ordinary skill in the art would reasonably doubt the asserted utility." *In re Brana*, 51 F.3d 1560, 1566, 34 U.S.P.Q.2d 1436 (Fed. Cir. 1995). Only then does the burden shift to the Applicant to provide rebuttal evidence. *Id.* As stated in the M.P.E.P., such rebuttal evidence does not need to absolutely prove that the asserted utility is real. Rather, the evidence only needs to be reasonably indicative of the asserted utility.

Applicants remind the PTO that the M.P.E.P. cautions that rejections for lack of utility are rarely sustained by federal courts, and that generally speaking, a utility rejection was

Appl. No. : 10/063,599
Filed : May 3, 2002

sustained because the applicant asserted a utility “that could **only be true if it violated a scientific principle, such as the second law of thermodynamics, or a law of nature, or was wholly inconsistent with contemporary knowledge in the art.**” M.P.E.P. § 2107.02 III B., citing *In re Gazave*, 379 F.2d 973, 978, 154 U.S.P.Q. 92, 96 (CCPA 1967) (underline emphasis in original, bold emphasis added). Rather than being wholly inconsistent with contemporary knowledge in the art, Applicants’ asserted utility is squarely within the teaching of leading textbooks in the field, and is supported by numerous references and the declarations of skilled experts.

Applicants’ asserted utility is based on the assertion that changes in mRNA level generally result in corresponding changes in the level of the encoded protein. In rejecting this conclusion, the PTO has cited references by Hu *et al.*, Haynes *et al.*, Gygi *et al.*, Anderson *et al.*, Lian *et al.*, and Fessler *et al.*

As explained above, these references are largely irrelevant when determining whether Applicants’ asserted utility is more likely than not true. Given the lack of support for the PTO’s position, Applicants submit that the PTO has not met its initial burden of overcoming the presumption that the asserted utility is sufficient to satisfy the utility requirement. And even if the PTO has met that burden, the Applicants’ supporting rebuttal evidence, including three uncontested expert declarations, excerpts from three textbooks, and over 115 scientific articles, is more than sufficient to establish that one of skill in the art would be more likely than not to believe that the claimed antibodies can be used as diagnostic tools for cancer, particularly esophageal, stomach, lung, rectal and melanoma tumor.

Specific Utility

The Asserted Substantial Utilities are Specific to the Claimed Antibodies

Applicants next address the PTO’s assertion that the asserted utilities are not specific to the claimed PRO1327 antibodies. Applicants respectfully disagree.

Specific utility is defined as utility which is “specific to the subject matter claimed,” in contrast to “a general utility that would be applicable to the broad class of the invention.” M.P.E.P. § 2107.01 I. Applicants submit that the evidence of differential expression of the PRO1327 gene and polypeptide in esophageal, stomach, lung, rectal and melanoma tumor cells,

Appl. No. : 10/063,599
Filed : May 3, 2002

along with the declarations and references discussed above, provide a specific utility for the claimed antibodies.

As discussed above, there are significant data which show that the gene for the PRO1327 polypeptide is differentially expressed in esophageal, stomach, lung, rectal and melanoma tumor tissue compared to normal tissue of these same types. These data are strong evidence that the PRO1327 gene and polypeptide are associated with esophageal, stomach, lung, rectal and melanoma tumors. Thus, contrary to the assertions of the PTO, Applicants submit that they have provided evidence associating the PRO1327 gene and polypeptide with specific diseases. The asserted utility as a diagnostic tool for cancer, particularly esophageal, stomach, lung, rectal and melanoma tumor, is a specific utility – it is not a general utility that would apply to the broad class of antibodies.

Conclusion

The PTO has asserted that the state of the art is such that polypeptide levels cannot be accurately predicted from mRNA levels. Applicants have addressed each of the PTO's supporting references and shown that they are either irrelevant, or taken as a whole, actually support Applicants' assertion that a change in mRNA level leads to a corresponding change in the level of the encoded protein. In addition, Applicants have submitted expert declarations, textbook excerpts, and over 115 scientific publications which support Applicants' asserted utility.

Given the totality of the evidence provided, Applicants submit that they have established a substantial, specific, and credible utility for the claimed antibodies as diagnostic tools. According to the PTO Utility Examination Guidelines (2001), irrefutable proof of a claimed utility is not required. Rather, a specific, substantial, and credible utility requires only a "reasonable" confirmation of a real world context of use. Applicants remind the PTO that:

A small degree of utility is sufficient . . . The claimed invention must only be capable of performing **some** beneficial function . . . An invention does not lack utility merely because the particular embodiment disclosed in the patent lacks perfection or performs crudely... A commercially successful product is not required... Nor is it essential that the invention accomplish all its intended functions... or operate under all conditions... partial success being sufficient to demonstrate patentable utility... In short, **the defense of non-utility cannot be sustained without proof of total incapacity**. If an invention is only partially

Appl. No. : **10/063,599**
Filed : **May 3, 2002**

successful in achieving a useful result, a rejection of the claimed invention as a whole based on a lack of utility is not appropriate. M.P.E.P. at 2107.01 (underline emphasis in original, bold emphasis added, citations omitted).

Applicants submit that they have established that it is more likely than not that one of skill in the art would reasonably accept the utility for the claimed PRO1327 antibodies set forth in the specification. In view of the above, Applicants respectfully request that the PTO reconsider and withdraw the utility rejection under 35 U.S.C. §101.

Rejections under 35 U.S.C. § 112, first paragraph – Enablement

The PTO maintains its rejection of Claims 1-5 as lacking enablement. The PTO states that because the claimed invention is not supported by either a specific and substantial asserted utility or a well established utility, one skilled in the art would not know how to use the claimed invention.

Applicants submit that in the discussion of the 35 U.S.C. § 101 rejection above, Applicants have established a substantial, specific, and credible utility for the claimed antibodies. Applicants respectfully request that to the extent the enablement rejection is based on a lack of utility, the PTO reconsider and withdraw the enablement rejection under 35 U.S.C. §112.

CONCLUSION

In view of the above, Applicants respectfully maintain that claims are patentable and request that they be passed to issue. Applicants invite the Examiner to call the undersigned if any remaining issues may be resolved by telephone.


Appl. No. : 10/063,599
Filed : May 3, 2002

Please charge any additional fees, including any fees for additional extension of time, or credit overpayment to Deposit Account No. 11-1410.

Respectfully submitted,

KNOBBE, MARTENS, OLSON & BEAR, LLP

Dated: June 14, 2006


By: 
AnneMarie Kaiser
Registration No. 37,649
Attorney of Record
Customer No. 30,313
(619) 235-8550

2552030
042506

EXHIBIT 1

Proteomics. 2005 Mar;5(4):894-906.

[Related Articles, Links](#)



A transcriptomic and proteomic analysis of the effect of CpG-ODN on human THP-1 monocytic leukemia cells.

Kuo CC, Kuo CW, Liang CM, Liang SM.

Institute of BioAgricultural Sciences, Academia Sinica, Taipei, Taiwan.

The CpG motif of bacterial DNA (CpG-DNA) is a potent immunostimulating agent whose mechanism of action is not yet clear. Here, we used both DNA microarray and proteomic approaches to investigate the effects of oligodeoxynucleotides containing the CpG motif (CpG-ODN) on gene transcription and protein expression profiles of CpG-ODN responsive THP-1 cells. Microarray analysis revealed that 2 h stimulation with CpG-ODN up-regulated 50 genes and down-regulated five genes. These genes were identified as being associated with inflammation, antimicrobial defense, transcriptional regulation, signal transduction, tumor progression, cell differentiation, proteolysis and metabolism. Longer stimulation (8 h) with CpG-ODN enhanced transcriptional expression of 58 genes. Among these 58 genes, none except one, namely WNT1 inducible signaling pathway protein 2, was the same as those induced after 2 h stimulation. Proteomic analysis by two-dimensional gel electrophoresis, followed by mass spectrometry identified several proteins up-regulated by CpG-ODN. These proteins included heat shock proteins, modulators of inflammation, metabolic proteins and energy pathway proteins. Comparison of microarray and proteomic expression profiles showed poor correlation. Use of more reliable and sensitive analyses, such as reverse transcriptase polymerase chain reaction, Western blotting and functional assays, on several genes and proteins, nonetheless, confirmed that there is indeed good correlation between mRNA and protein expression after CpG-ODN treatment. This study also revealed that several anti-apoptotic and neuroprotective related proteins, not previously reported, are activated by CpG-DNA. These findings have extended our knowledge on the activation of cells by CpG-DNA and may contribute to further understanding of mechanisms that link innate immunity with acquired immune response(s).

PMID: 15693060 [PubMed - indexed for MEDLINE]

Genome-wide Study of Gene Copy Numbers, Transcripts, and Protein Levels in Pairs of Non-invasive and Invasive Human Transitional Cell Carcinomas*

Torben F. Ørntoft§, Thomas Thykjaer||, Frederic M. Waldman||, Hans Wolf**, and Julo E. Celis††

Gain and loss of chromosomal material is characteristic of bladder cancer, as well as malignant transformation in general. The consequences of these changes at both the transcription and translation levels is at present unknown partly because of technical limitations. Here we have attempted to address this question in pairs of non-invasive and invasive human bladder tumors using a combination of technology that included comparative genomic hybridization, high density oligonucleotide array-based monitoring of transcript levels (5600 genes), and high resolution two-dimensional gel electrophoresis. The results showed that there is a gene dosage effect that in some cases superimposes on other regulatory mechanisms. This effect depended ($p < 0.015$) on the magnitude of the comparative genomic hybridization change. In general (18 of 23 cases), chromosomal areas with more than 2-fold gain of DNA showed a corresponding increase in mRNA transcripts. Areas with loss of DNA, on the other hand, showed either reduced or unaltered transcript levels. Because most proteins resolved by two-dimensional gels are unknown it was only possible to compare mRNA and protein alterations in relatively few cases of well focused abundant proteins. With few exceptions we found a good correlation ($p < 0.005$) between transcript alterations and protein levels. The implications, as well as limitations, of the approach are discussed. *Molecular & Cellular Proteomics* 1:37-45, 2002.

Aneuploidy is a common feature of most human cancers (1), but little is known about the genome-wide effect of this

From the §Department of Clinical Biochemistry, Molecular Diagnostic Laboratory and **Department of Urology, Aarhus University Hospital, Skejby, DK-8200 Aarhus N, Denmark, ||AROS Applied Biotechnology ApS, Gustav Wiedsvej 10, DK-8000 Aarhus C, Denmark, ||UCSF Cancer Center and Department of Laboratory Medicine, University of California, San Francisco, CA 94143-0808, and ††Institute of Medical Biochemistry and Danish Centre for Human Genome Research, Ole Worms Allé 170, Aarhus University, DK-8000 Aarhus C, Denmark

Received, September 26, 2001, and in revised form, November 7, 2001

Published, MCP Papers in Press, November 13, 2001, DOI 10.1074/mcp.M100019-MCP200

phenomenon at both the transcription and translation levels. High throughput array studies of the breast cancer cell line BT474 has suggested that there is a correlation between DNA copy numbers and gene expression in highly amplified areas (2), and studies of individual genes in solid tumors have revealed a good correlation between gene dose and mRNA or protein levels in the case of c-erb-B2, cyclin d1, *ems1*, and N-myc (3-5). However, a high cyclin D1 protein expression has been observed without simultaneous amplification (4), and a low level of c-myc copy number increase was observed without concomitant c-myc protein overexpression (6).

In human bladder tumors, karyotyping, fluorescent *in situ* hybridization, and comparative genomic hybridization (CGH)¹ have revealed chromosomal aberrations that seem to be characteristic of certain stages of disease progression. In the case of non-invasive pTa transitional cell carcinomas (TCCs), this includes loss of chromosome 9 or parts of it, as well as loss of Y in males. In minimally invasive pT1 TCCs, the following alterations have been reported: 2q-, 11p-, 1q+, 11q13+, 17q+, and 20q+ (7-12). It has been suggested that these regions harbor tumor suppressor genes and oncogenes; however, the large chromosomal areas involved often contain many genes, making meaningful predictions of the functional consequences of losses and gains very difficult.

In this investigation we have combined genome-wide technology for detecting genomic gains and losses (CGH) with gene expression profiling techniques (microarrays and proteomics) to determine the effect of gene copy number on transcript and protein levels in pairs of non-invasive and invasive human bladder TCCs.

EXPERIMENTAL PROCEDURES

Material—Bladder tumor biopsies were sampled after informed consent was obtained and after removal of tissue for routine pathology examination. By light microscopy tumors 335 and 532 were staged by an experienced pathologist as pTa (superficial papillary).

¹ The abbreviations used are: CGH, comparative genomic hybridization; TCC, transitional cell carcinoma; LOH, loss of heterozygosity; PA-FABP, psoriasis-associated fatty acid-binding protein; 2D, two-dimensional.

Gene Copy Numbers, Transcripts, and Protein Levels




FIG. 1. DNA copy number and mRNA expression level. Shown from left to right are chromosome (Chr.), CGH profiles, gene location and expression level of specific genes, and overall expression level along the chromosome. **A**, expression of mRNA in invasive tumor 733 as compared with the non-invasive counterpart tumor 335. The average fluorescent signal ratio between tumor DNA and normal DNA is shown along the length of the chromosome (left). The bold curve in the ratio profile represents a mean of four chromosomes and is surrounded by (thin curves) indicating one standard deviation. The central vertical line (broken) indicates a ratio value of 1 (no change), and the vertical lines next to it (dotted) indicate a ratio of 0.5 (left) and 2.0 (right). In chromosomes where the non-invasive tumor 335 used for comparison showed alterations in DNA content, the ratio profile of that chromosome is shown to the right of the invasive tumor profile. The colored bars represent one gene each, identified by the running numbers above the bars (the name of the gene can be seen at www.MDL.DK/sdata.html). The bars indicate the purported location of the gene, and the colors indicate the expression level of the gene in the invasive tumor compared with the non-invasive counterpart; >2-fold increase (black), >2-fold decrease (blue), no significant change (orange). The bar to the far right, entitled Expression shows the resulting change in expression along the chromosome; the colors indicate that at least half of the genes were up-regulated (black), at least half of the genes down-regulated (blue), or more than half of the genes are unchanged (orange). If a gene was absent in one of the samples and present in another, it was regarded as more than a 2-fold change. A 2-fold level was chosen as this corresponded to one standard deviation in a double determination of ~1800 genes. Centromeres and heterochromatic regions were excluded from data analysis.

grade I and II, respectively, tumors 733 and 827 were staged as pT1 (invasive into submucosa), 733 was staged as solid, and 827 was staged as papillary, both grade III.

mRNA Preparation—Tissue biopsies, obtained fresh from surgery, were embedded immediately in a sodium-guanidinium thiocyanate solution and stored at -80 °C. Total RNA was isolated using the RNeasy B RNA isolation method (WAK-Chemie Medical GmbH). poly(A)⁺ RNA was isolated by an oligo(dT) selection step (Oligotex mRNA kit; Qiagen).

cRNA Preparation—1 µg of mRNA was used as starting material. The first and second strand cDNA synthesis was performed using the SuperScript[®] choice system (Invitrogen) according to the manufacturer's instructions but using an oligo(dT) primer containing a T7 RNA polymerase binding site. Labeled cRNA was prepared using the ME-GAscript[®] *in vitro* transcription kit (Ambion). Biotin-labeled CTP and

UTP (Enzo) was used, together with unlabeled NTPs in the reaction. Following the *in vitro* transcription reaction, the unincorporated nucleotides were removed using RNeasy columns (Qiagen).

Array Hybridization and Scanning—Array hybridization and scanning was modified from a previous method (13). 10 µg of cRNA was fragmented at 94 °C for 35 min in buffer containing 40 mM Tris acetate, pH 8.1, 100 mM KOAc, 30 mM MgOAc. Prior to hybridization, the fragmented cRNA in a 6× SSPE-T hybridization buffer (1 M NaCl, 10 mM Tris, pH 7.8, 0.005% Triton), was heated to 85 °C for 5 min, subsequently cooled to 40 °C, and loaded onto the Affymetrix probe array cartridge. The probe array was then incubated for 16 h at 40 °C at constant rotation (60 rpm). The probe array was exposed to 10 washes in 6× SSPE-T at 25 °C followed by 4 washes in 0.5× SSPE-T at 50 °C. The biotinylated cRNA was stained with a streptavidin-phycoerythrin conjugate, 10 µg/ml (Molecular Probes) in 6× SSPE-T




Fig. 1—continued

for 30 min at 25 °C followed by 10 washes in 6× SSPE-T at 25 °C. The probe arrays were scanned at 560 nm using a confocal laser scanning microscope (made for Affymetrix by Hewlett-Packard). The readings from the quantitative scanning were analyzed by Affymetrix gene expression analysis software.

Microsatellite Analysis—Microsatellite Analysis was performed as described previously (14). Microsatellites were selected by use of www.ncbi.nlm.nih.gov/genemap98, and primer sequences were obtained from the genome data base at www.gdb.org. DNA was extracted from tumor and blood and amplified by PCR in a volume of 20 μ l for 35 cycles. The amplicons were denatured and electrophoresed for 3 h in an ABI Prism 377. Data were collected in the Gene Scan program for fragment analysis. Loss of heterozygosity was defined as less than 33% of one allele detected in tumor amplicons compared with blood.

Proteomic Analysis—TCCs were minced into small pieces and homogenized in a small glass homogenizer in 0.5 ml of lysis solution. Samples were stored at -20 °C until use. The procedure for 2D gel electrophoresis has been described in detail elsewhere (15, 16). Gels were stained with silver nitrate and/or Coomassie Brilliant Blue. Proteins were identified by a combination of procedures that included microsequencing, mass spectrometry, two-dimensional gel Western immunoblotting, and comparison with the master two-dimensional gel image of human keratinocyte proteins; see blobase.dk/cgi-bin/cells.

CGH—Hybridization of differentially labeled tumor and normal DNA to normal metaphase chromosomes was performed as described previously (10). Fluorescein-labeled tumor DNA (200 ng), Texas Red-

labeled reference DNA (200 ng), and human Cot-1 DNA (20 μ g) were denatured at 37 °C for 5 min and applied to denatured normal metaphase slides. Hybridization was at 37 °C for 2 days. After washing, the slides were counterstained with 0.15 μ g/ml 4,6-diamidino-2-phenylindole in an anti-fade solution. A second hybridization was performed for all tumor samples using fluorescein-labeled reference DNA and Texas Red-labeled tumor DNA (inverse labeling) to confirm the aberrations detected during the initial hybridization. Each CGH experiment also included a normal control hybridization using fluorescein- and Texas Red-labeled normal DNA. Digital image analysis was used to identify chromosomal regions with abnormal fluorescence ratios, indicating regions of DNA gains and losses. The average green:red fluorescence intensity ratio profiles were calculated using four images of each chromosome (eight chromosomes total) with normalization of the green:red fluorescence intensity ratio for the entire metaphase and background correction. Chromosome identification was performed based on 4,6-diamidino-2-phenylindole banding patterns. Only images showing uniform high intensity fluorescence with minimal background staining were analyzed. All centromeres, p arms of acrocentric chromosomes, and heterochromatic regions were excluded from the analysis.

RESULTS

Comparative Genomic Hybridization—The CGH analysis identified a number of chromosomal gains and losses in the

Gene Copy Numbers, Transcripts, and Protein Levels

TABLE I
Correlation between alterations detected by CGH and by expression monitoring

Top, CGH used as independent variable (if CGH alteration – what expression ratio was found); bottom, altered expression used as independent variable (if expression alteration – what CGH deviation was found).

CGH alterations	Tumor 733 vs. 335		Concordance	CGH alterations	Tumor 827 vs. 532		Concordance
	Expression change clusters				Expression change clusters		
13 Gain	10 Up-regulation 0 Down-regulation 3 No change	77%		10 Gain	8 Up-regulation 0 Down-regulation 2 No change	80%	
10 Loss	1 Up-regulation 6 Down-regulation 4 No change	50%		12 Loss	3 Up-regulation 2 Down-regulation 7 No change	17%	
Expression change clusters	Tumor 733 vs. 335		Concordance	Expression change clusters	Tumor 827 vs. 532		Concordance
	CGH alterations				CGH alterations		
16 Up-regulation	11 Gain 2 Loss 3 No change	69%		17 Up-regulation	10 Gain 5 Loss 2 No change	59%	
21 Down-regulation	1 Gain 8 Loss 12 No change	38%		9 Down-regulation	0 Gain 9 Loss 6 No change	33%	
15 No change	3 Gain 3 Loss 9 No change	60%		21 No change	1 Gain 3 Loss 17 No change	81%	

two invasive tumors (stage pT1, TCCs 733 and 827), whereas the two non-invasive papillomas (stage pTa, TCCs 335 and 532) showed only 9p–, 9q22–q33–, and X–, and 7+, 9q–, and Y–, respectively. Both invasive tumors showed changes (1q22–24+, 2q14.1–qter–, 3q12–q13.3–, 6q12–q22–, 9q34+, 11q12–q13+, 17+, and 20q11.2–q12+) that are typical for their disease stage, as well as additional alterations, some of which are shown in Fig. 1. Areas with gains and losses deviated from the normal copy number to some extent, and the average numerical deviation from normal was 0.4-fold in the case of TCC 733 and 0.3-fold for TCC 827. The largest changes, amounting to at least a doubling of chromosomal content, were observed at 1q23 in TCC 733 (Fig. 1A) and 20q12 in TCC 827 (Fig. 1B).

mRNA Expression in Relation to DNA Copy Number—The mRNA levels from the two invasive tumors (TCCs 827 and 733) were compared with the two non-invasive counterparts (TCCs 532 and 335). This was done in two separate experiments in which we compared TCCs 733 to 335 and 827 to 532, respectively, using two different scaling settings for the arrays to rule out scaling as a confounding parameter. Approximately 1,800 genes that yielded a signal on the arrays were searched in the Unigene and Genemap data bases for chromosomal location, and those with a known location (1096) were plotted as bars covering their purported locus. In that way it was possible to construct a graphic presentation of DNA copy number and relative mRNA levels along the individual chromosomes (Fig. 1).

For each mRNA a ratio was calculated between the level in the invasive versus the non-invasive counterpart. Bars, which represent chromosomal location of a gene, were color-coded according to the expression ratio, and only differences larger

than 2-fold were regarded as informative (Fig. 1). The density of genes along the chromosomes varied, and areas containing only one gene were excluded from the calculations. The resolution of the CGH method is very low, and some of the outlier data may be because of the fact that the boundaries of the chromosomal aberrations are not known at high resolution.

Two sets of calculations were made from the data. For the first set we used CGH alterations as the independent variable and estimated the frequency of expression alterations in these chromosomal areas. In general, areas with a strong gain of chromosomal material contained a cluster of genes having increased mRNA expression. For example, both chromosomes 1q21–q25, 2p and 9q, showed a relative gain of more than 100% in DNA copy number that was accompanied by increased mRNA expression levels in the two tumor pairs (Fig. 1). In most cases, chromosomal gains detected by CGH were accompanied by an increased level of transcripts in both TCCs 733 (77%) and 827 (80%) (Table I, top). Chromosomal losses, on the other hand, were not accompanied by decreased expression in several cases, and were often registered as having unaltered RNA levels (Table I, top). The inability to detect RNA expression changes in these cases was not because of fewer genes mapping to the lost regions (data not shown).

In the second set of calculations we selected expression alterations above 2-fold as the independent variable and estimated the frequency of CGH alterations in these areas. As above, we found that increased transcript expression correlated with gain of chromosomal material (TCC 733, 69% and TCC 827, 59%), whereas reduced expression was often detected in areas with unaltered CGH ratios (Table I, bottom). Furthermore, as a control we looked at areas with no alter-




Fig. 2. Correlation between maximum CGH aberration and the ability to detect expression change by oligonucleotide array monitoring. The aberration is shown as a numerical -fold change in ratio between invasive tumors 827 (▲) and 733 (◆) and their non-invasive counterparts 532 and 335. The expression change was taken from the *Expressari* line to the right in Fig. 1, which depicts the resulting expression change for a given chromosomal region. At least half of the mRNAs from a given region have to be either up- or down-regulated to be scored as an expression change. All chromosomal arms in which the CGH ratio plus or minus one standard deviation was outside the ratio value of one were included.

ation in expression. No alteration was detected by CGH in most of these areas (TCC 733, 60% and TCC 827, 81%; see Table I, bottom). Because the ability to observe reduced or increased mRNA expression clustering to a certain chromosomal area clearly reflected the extent of copy number changes, we plotted the maximum CGH aberrations in the regions showing CGH changes against the ability to detect a change in mRNA expression as monitored by the oligonucleotide arrays (Fig. 2). For both tumors TCC 733 ($p < 0.015$) and TCC 827 ($p < 0.00003$) a highly significant correlation was observed between the level of CGH ratio change (reflecting the DNA copy number) and alterations detected by the array based technology (Fig. 2). Similar data were obtained when areas with altered expression were used as independent variables. These areas correlated best with CGH when the CGH ratio deviated 1.6- to 2.0-fold (Table I, bottom) but mostly did not at lower CGH deviations. These data probably reflect that loss of an allele may only lead to a 50% reduction in expression level, which is at the cut-off point for detection of expression alterations. Gain of chromosomal material can occur to a much larger extent.

Microsatellite-based Detection of Minor Areas of Losses—In TCC 733, several chromosomal areas exhibiting DNA amplification were preceded or followed by areas with a normal CGH but reduced mRNA expression (see Fig. 1, TCC 733 chromosome 1q32, 2p21, and 7q21 and q32, 9q34, and 10q22). To determine whether these results were because of undetected loss of chromosomal material in these regions or

because of other non-structural mechanisms regulating transcription, we examined two microsatellites positioned at chromosome 1q25–32 and two at chromosome 2p22. Loss of heterozygosity (LOH) was found at both 1q25 and at 2p22 indicating that minor deleted areas were not detected with the resolution of CGH (Fig. 3). Additionally, chromosome 2p in TCC 733 showed a CGH pattern of gain/no change/gain of DNA that correlated with transcript increase/decrease/increase. Thus, for the areas showing increased expression there was a correlation with the DNA copy number alterations (Fig. 1A). As indicated above, the mRNA decrease observed in the middle of the chromosomal gain was because of LOH, implying that one of the mechanisms for mRNA down-regulation may be regions that have undergone smaller losses of chromosomal material. However, this cannot be detected with the resolution of the CGH method.

In both TCC 733 and TCC 827, the telomeric end of chromosome 11p showed a normal ratio in the CGH analysis; however, clusters of five and three genes, respectively, lost their expression. Two microsatellites (D11S1760, D11S922) positioned close to MUC2, IGF2, and cathepsin D indicated LOH as the most likely mechanism behind the loss of expression (data not shown).

A reduced expression of mRNA observed in TCC 733 at chromosomes 3q24, 11p11, 12p12.2, 12q21.1, and 16q24 and in TCC 827 at chromosome 11p15.5, 12p11, 15q11.2, and 18q12 was also examined for chromosomal losses using microsatellites positioned as close as possible to the gene loci




FIG. 3. Microsatellite analysis of loss of heterozygosity. Tumor 733 showing loss of heterozygosity at chromosome 1q25, detected (a) by D1S215 close to HLA class I histocompatibility antigen (gene number 38 in Fig. 1), (b) by D1S2735 close to cathepsin E (gene number 41 in Fig. 1), and (c) at chromosome 2p23 by D2S2251 close to general β -spectrin (gene number 11 on Fig. 1) and of (d) tumor 827 showing loss of heterozygosity at chromosome 18q12 by S18S1118 close to mitochondrial 3-oxoacyl-coenzyme A thiolase (gene number 12 in Fig. 1). The upper curves show the electropherogram obtained from normal DNA from leukocytes (N), and the lower curves show the electropherogram from tumor DNA (T). In all cases one allele is partially lost in the tumor amplicon.

showing reduced mRNA transcripts. Only the microsatellite positioned at 18q12 showed LOH (Fig. 3), suggesting that transcriptional down-regulation of genes in the other regions may be controlled by other mechanisms.

Relation between Changes in mRNA and Protein Levels—2D-PAGE analysis, in combination with Coomassie Brilliant Blue and/or silver staining, was carried out on all four tumors using fresh biopsy material. 40 well resolved abundant known proteins migrating in areas away from the edges of the pH




FIG. 4. Correlation between protein levels as judged by 2D-PAGE and transcript ratio. For comparison proteins were divided in three groups, unaltered in level or up- or down-regulated (horizontal axis). The mRNA ratio as determined by oligonucleotide arrays was plotted for each gene (vertical axis). Δ , mRNAs that were scored as present in both tumors used for the ratio calculation; ∇ , mRNAs that were scored as absent in the invasive tumors (along horizontal axis) or as absent in non-invasive reference (top of figure). Two different scalings were used to exclude scaling as a confounder, TCCs 827 and 532 (Δ) were scaled with background suppression, and TCCs 733 and 335 (∇) were scaled without suppression. Both comparisons showed highly significant ($p < 0.005$) differences in mRNA ratios between the groups. Proteins shown were as follows: Group A (from left), phosphoglucomutase 1, glutathione transferase class μ number 4, fatty acid-binding protein homologue, cytochrome P-450, keratin 13; B (from left), fatty acid-binding protein homologue, 28-kDa heat shock protein, cytochrome P-450, and calyculin; C (from left), α -enolase, hnRNP B1, 28-kDa heat shock protein, 14-3-3- ϵ , and pre-mRNA splicing factor; D, mesothelial keratin K7 (type II); E (from top), glutathione S-transferase- π and mesothelial keratin K7 (type II); F (from top and left), adenylyl cyclase-associated protein, E-cadherin, keratin 19, calgizzarin, phosphoglycerate mutase, annexin IV, cytoskeletal γ -actin, hnRNP A1, integral membrane protein calnexin (IP90), hnRNP H, brain-type clathrin light chain- α , hnRNP F, 70-kDa heat shock protein, heterogeneous nuclear ribonucleoprotein A/B, translationally controlled tumor protein, liver glyceraldehyde-3-phosphate dehydrogenase, keratin 8, aldehyde reductase, and Na,K-ATPase β -1 subunit; G, (from top and left), TCP20, calgizzarin, 70-kDa heat shock protein, calnexin, hnRNP H, cytochrome P-450, ATP synthase, keratin 19, triosephosphate isomerase, hnRNP F, liver glyceraldehyde-3-phosphate dehydrogenase, glutathione S-transferase- π , and keratin 8; H (from left), plasma gelsolin, autoantigen calreticulin, thioredoxin, and NAD $^{+}$ -dependent 15-hydroxyprostaglandin dehydrogenase; I (from top), prollyl 4-hydroxylase β -subunit, cytochrome P-450, cytochrome P-17, prohibitin, and fructose 1,6-bisphosphatase; J annexin II; K, annexin IV; L (from top and left), 90-kDa heat shock protein, prollyl 4-hydroxylase β -subunit, α -enolase, GRP 78, cyclophilin, and cofillin.

gradient, and having a known chromosomal location, were selected for analysis in the TCC pair 827/532. Proteins were identified by a combination of methods (see "Experimental Procedures"). In general there was a highly significant correlation ($p < 0.005$) between mRNA and protein alterations (Fig. 4). Only one gene showed disagreement between transcript alteration and protein alteration. Except for a group of cyto-




Fig. 5. Comparison of protein and transcript levels in invasive and non-invasive TCCs. The upper part of the figure shows a 2D gel (left) and the oligonucleotide array (right) of TCC 532. The red rectangles on the upper gel highlight the areas that are compared below. Identical areas of 2D gels of TCCs 532 and 827 are shown below. Clearly, cytokeratins 13 and 15 are strongly down-regulated in TCC 827 (red annotation). The tile on the array containing probes for cytokeratin 15 is enlarged below the array (red arrow) from TCC 532 and is compared with TCC 827. The upper row of squares in each tile corresponds to perfect match probes; the lower row corresponds to mismatch probes containing a mutation (used for correction for unspecific binding). Absence of signal is depicted as black, and the higher the signal the lighter the color. A high transcript level was detected in TCC 532 (6151 units) whereas a much lower level was detected in TCC 827 (absence of signals). For cytokeratin 13, a high transcript level was also present in TCC 532 (15859 units), and a much lower level was present in TCC 827 (623 units). The 2D gels at the bottom of the figure (left) show levels of PA-FABP and adipocyte-FABP in TCCs 335 and 733 (invasive), respectively. Both proteins are down-regulated in the invasive tumor. To the right we show the array tiles for the PA-FABP transcript. A medium transcript level was detected in the case of TCC 335 (1277 units) whereas very low levels were detected in TCC 733 (166 units). IEF, Isoelectric focusing.

keratins encoded by genes on chromosome 17 (Fig. 5) the analyzed proteins did not belong to a particular family. 26 well focused proteins whose genes had a known chromosomal location were detected in TCCs 733 and 335, and of these 19 correlated ($p < 0.005$) with the mRNA changes detected using the arrays (Fig. 4). For example, PA-FABP was highly expressed in the non-invasive TCC 335 but lost in the invasive counterpart (TCC 733; see Fig. 5). The smaller number of proteins detected in both 733 and 335 was because of the smaller size of the biopsies that were available.

11 chromosomal regions where CGH showed aberrations that corresponded to the changes in transcript levels also showed corresponding changes in the protein level (Table II). These regions included genes that encode proteins that are found to be frequently altered in bladder cancer, namely cytokeratins 17 and 20, annexins II and IV, and the fatty acid-binding proteins PA-FABP and FABP1. Four of these proteins were encoded by genes in chromosome 17q, a frequently amplified chromosomal area in invasive bladder cancers.

DISCUSSION

Most human cancers have abnormal DNA content, having lost some chromosomal parts and gained others. The present study provides some evidence as to the effect of these gains and losses on gene expression in two pairs of non-invasive and invasive TCCs using high throughput expression arrays and proteomics, in combination with CGH. In general, the results showed that there is a clear individual regulation of the mRNA expression of single genes, which in some cases was superimposed by a DNA copy number effect. In most cases, genes located in chromosomal areas with gains often exhibited increased mRNA expression, whereas areas showing losses showed either no change or a reduced mRNA expression. The latter might be because of the fact that losses most often are restricted to loss of one allele, and the cut-off point for detection of expression alterations was a 2-fold change, thus being at the border of detection. In several cases, how-

TABLE II
Proteins whose expression level correlates with both mRNA and gene dose changes

Protein	Chromosomal location	Tumor TCC	CGH alteration	Transcript alteration ^a	Protein alteration
Annexin II	1q21	733	Gain	Abs to Pres ^a	Increase
Annexin IV	2p13	733	Gain	3.9-Fold up	Increase
Cytokeratin 17	17q12-q21	827	Gain	3.8-Fold up	Increase
Cytokeratin 20	17q21.1	827	Gain	5.8-Fold up	Increase
(PA-)FABP	8q21.2	827	Loss	10-Fold down	Decrease
FABP1	9q22	827	Gain	2.3-Fold up	Increase
Plasma gelsolin	9q31	827	Gain	Abs to Pres	Increase
Heat shock protein 28	15q12-q13	827	Loss	2.5-Fold up	Decrease
Prohibitin	17q21	827/733	Gain	3.7-/2.5-Fold up ^b	Increase
Prolyl-4-hydroxyl	17q25	827/733	Gain	6.7-/1.6-Fold up	Increase
hnRNPB1	7p15	827	Loss	2.5-Fold down	Decrease

^a Abs, absent; Pres, present.

^b In cases where the corresponding alterations were found in both TCCs 827 and 733 these are shown as 827/733.

ever, an increase or decrease in DNA copy number was associated with *de novo* occurrence or complete loss of transcript, respectively. Some of these transcripts could not be detected in the non-invasive tumor but were present at relatively high levels in areas with DNA amplifications in the invasive tumors (e.g. In TCC 733 transcript from cellular ligand of annexin II gene (chromosome 1q21) from absent to 2670 arbitrary units; In TCC 827 transcript from small proline-rich protein 1 gene (chromosome 1q12-q21.1) from absent to 1326 arbitrary units). It may be anticipated from these data that significant clustering of genes with an increased expression to a certain chromosomal area indicates an increased likelihood of gain of chromosomal material in this area.

Considering the many possible regulatory mechanisms acting at the level of transcription, it seems striking that the gene dose effects were so clearly detectable in gained areas. One hypothetical explanation may lie in the loss of controlled methylation in tumor cells (17-19). Thus, it may be possible that in chromosomes with increased DNA copy numbers two or more alleles could be demethylated simultaneously leading to a higher transcription level, whereas in chromosomes with losses the remaining allele could be partly methylated, turning off the process (20, 21). A recent report has documented a ploidy regulation of gene expression in yeast, but in this case all the genes were present in the same ratio (22), a situation that is not analogous to that of cancer cells, which show marked chromosomal aberrations, as well as gene dosage effects.

Several CGH studies of bladder cancer have shown that some chromosomal aberrations are common at certain stages of disease progression, often occurring in more than 1 of 3 tumors. In pTa tumors, these include 9p-, 9q-, 1q+, Y- (2, 6), and in pT1 tumors, 2q-, 11p-, 11q-, 1q+, 5p+, 8q+, 17q+, and 20q+ (2-4, 6, 7). The pTa tumors studied here showed similar aberrations such as 9p- and 9q22-q33- and 9q- and Y-, respectively. Likewise, the two minimal invasive pT1 tumors showed aberrations that are commonly seen at that stage, and TCC 827 had a remarkable resemblance to the commonly seen pattern of losses and gains, such as 1q22-24 amplification (seen in both tumors), 11q14-q22 loss, the latter often linked to 17q+ (both tumors), and 1q+ and 9p-, often linked to 20q+ and 11q13+ (both tumors) (7-9). These observations indicate that the pairs of tumors used in this study exhibit chromosomal changes observed in many tumors, and therefore the findings could be of general importance for bladder cancer.

Considering that the mapping resolution of CGH is of about 20 megabases it is only possible to get a crude picture of chromosomal instability using this technique. Occasionally, we observed reduced transcript levels close to or inside regions with increased copy numbers. Analysis of these regions by positioning heterozygous microsatellites as close as possible to the locus showing reduced gene expression revealed loss of heterozygosity in several cases. It seems likely that multiple and different events occur along each chromosomal

arm and that the use of cDNA microarrays for analysis of DNA copy number changes will reach a resolution that can resolve these changes, as has recently been proposed (2). The outlier data were not more frequent at the boundaries of the CGH aberrations. At present we do not know the mechanism behind chromosomal aneuploidy and cannot predict whether chromosomal gains will be transcribed to a larger extent than the two native alleles. A mechanism as genetic imprinting has an impact on the expression level in normal cells and is often reduced in tumors. However, the relation between imprinting and gain of chromosomal material is not known.

We regard it as a strength of this investigation that we were able to compare invasive tumors to benign tumors rather than to normal urothelium, as the tumors studied were biologically very close, and probably may represent successive steps in the progression of bladder cancer. Despite the limited amount of fresh tissue available it was possible to apply three different state of the art methods. The observed correlation between DNA copy number and mRNA expression is remarkable when one considers that different pieces of the tumor biopsies were used for the different sets of experiments. This indicates that bladder tumors are relatively homogenous, a notion recently supported by CGH and LOH data that showed a remarkable similarity even between tumors and distant metastasis (10, 23).

In the few cases analyzed, mRNA and protein levels showed a striking correspondence although in some cases we found discrepancies that may be attributed to translational regulation, post-translational processing, protein degradation, or a combination of these. Some transcripts belong to undertranslated mRNA pools, which are associated with few translationally inactive ribosomes; these pools, however, seem to be rare (24). Protein degradation, for example, may be very important in the case of polypeptides with a short half-life (e.g. signaling proteins). A poor correlation between mRNA and protein levels was found in liver cells as determined by arrays and 2D-PAGE (25), and a moderate correlation was recently reported by Ideker *et al.* (26) in yeast.

Interestingly, our study revealed a much better correlation between gained chromosomal areas and increased mRNA levels than between loss of chromosomal areas and reduced mRNA levels. In general, the level of CGH change determined the ability to detect a change in transcript. One possible explanation could be that by losing one allele the change in mRNA level is not so dramatic as compared with gain of material, which can be rather unlimited and may lead to a severalfold increase in gene copy number resulting in a much higher impact on transcript level. The latter would be much easier to detect on the expression arrays as the cut-off point was placed at a 2-fold level so as not to be biased by noise on the array. Construction of arrays with a better signal to noise ratio may in the future allow detection of lesser than 2-fold alterations in transcript levels, a feature that may facilitate the analysis of the effect of loss of chromosomal areas on transcript levels.

In eleven cases we found a significant correlation between DNA copy number, mRNA expression, and protein level. Four of these proteins were encoded by genes located at a frequently amplified area in chromosome 17q. Whether DNA copy number is one of the mechanisms behind alteration of these eleven proteins is at present unknown and will have to be proved by other methods using a larger number of samples. One factor making such studies complicated is the large extent of protein modification that occurs after translation, requiring immunoidentification and/or mass spectrometry to correctly identify the proteins in the gels.

In conclusion, the results presented in this study exemplify the large body of knowledge that may be possible to gather in the future by combining state of the art techniques that follow the pathway from DNA to protein (26). Here, we used a traditional chromosomal CGH method, but in the future high resolution CGH based on microarrays with many thousand radiation hybrid-mapped genes will increase the resolution and information derived from these types of experiments (2). Combined with expression arrays analyzing transcripts derived from genes with known locations, and 2D gel analysis to obtain information at the post-translational level, a clearer and more developed understanding of the tumor genome will be forthcoming.

Acknowledgments—We thank Mle Madsen, Hanne Steen, Inge Lis Thorsen, Hans Lund, Nikolaj Ørntoft, and Lynn Bjerke for technical help and Thomas Glingeras, Christine Harrington, and Morten Østergaard for valuable discussions.

*This work was supported by grants from The Danish Cancer Society, the University of Aarhus, Aarhus County, Novo Nordic, the Danish Biotechnology Program, the Frønkels Foundation, the John and Birthe Meyer Foundation, and NCI, National Institutes of Health Grant CA47537. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

§ To whom correspondence should be addressed: Dept. of Clinical Biochemistry, Molecular Diagnostic Laboratory, Aarhus University Hospital, Skøjby, DK-8200 Aarhus N, Denmark. Tel.: 45-89495100/45-86156201 (private); Fax: 45-89496018; E-mail: orntoft@kba.sks.au.dk


REFERENCES

1. Lengauer, C., Kinzler, K. W., and Vogelstein, B. (1998) Genetic instabilities in human cancers. *Nature* 396, 643-649.
2. Pollock, J. R., Perou, C. M., Alizadeh, A. A., Eisen, M. B., Pergamenschikov, A., Williams, C. F., Jeffrey, S. S., Botstein, D., and Brown, P. O. (1999) Genome-wide analysis of DNA copy-number changes using cDNA microarrays. *Nat. Genet.* 23, 41-46.
3. de Cremoux, P., Martin, E. C., Vincent-Salomon, A., Dieras, V., Barbaroux, C., Liva, S., Poullart, P., Sastre-Garau, X., and Magdelenat, H. (1999) Quantitative PCR analysis of c-erb B-2 (HER2/neu) gene amplification and comparison with p185(HER2/neu) protein expression in breast cancer drill biopsies. *Int. J. Cancer* 83, 157-161.
4. Brungler, P. P., Tamini, Y., Shuuring, E., and Schalken, J. (1996) Expression of cyclin D1 and EMS1 in bladder tumors; relationship with chromosome 11q13 amplifications. *Oncogene* 12, 1747-1753.
5. Slave, L., Eilenbogen, R., Jung, W. H., Vawter, G. F., Kretschmar, C., Grier, H., and Korf, B. R. (1990) myc gene amplification and expression in primary human neuroblastoma. *Cancer Res.* 50, 1459-1463.
6. Sauter, G., Carroll, P., Moch, H., Kallioniemi, A., Kerschmann, R., Narayan, P., Mihatsch, M. J., and Waldman, F. M. (1995) c-myc copy number gains in bladder cancer detected by fluorescence *in situ* hybridization. *Am. J. Pathol.* 148, 1131-1139.
7. Richter, J., Jlang, F., Gorog, J. P., Sarlantis, G., Egenter, C., Gasser, T. C., Moch, H., Mihatsch, M. J., and Sauter, G. (1997) Marked genetic differences between stage pTa and stage pT1 papillary bladder cancer detected by comparative genomic hybridization. *Cancer Res.* 57, 2860-2864.
8. Richter, J., Boffa, L., Wagner, U., Schraml, P., Gasser, T. C., Moch, H., Mihatsch, M. J., and Sauter, G. (1998) Patterns of chromosomal imbalances in advanced urinary bladder cancer detected by comparative genomic hybridization. *Am. J. Pathol.* 153, 1615-1621.
9. Bruch, J., Wotr, G., Hautmann, R., Mattfeldt, T., Bruderlein, S., Moller, P., Sauter, S., Hamelster, H., Vogel, W., and Palss, T. (1998) Chromosomal changes during progression of transitional cell carcinoma of the bladder and delineation of the amplified interval on chromosome arm 8q. *Genes Chromosomes Cancer* 23, 167-174.
10. Hovey, R. M., Chu, L., Balazs, M., De Vries, S., Moore, D., Sauter, G., Carroll, P. R., and Waldman, F. M. (1998) Genetic alterations in primary bladder cancers and their metastases. *Cancer Res.* 58, 3555-3560.
11. Simon, R., Burger, H., Brinkschmidt, C., Bookar, W., Hertle, L., and Terpe, H. J. (1998) Chromosomal aberrations associated with invasion in papillary superficial bladder cancer. *J. Pathol.* 185, 345-351.
12. Koo, S. H., Kwon, K. C., Ihm, C. H., Jeon, Y. M., Park, J. W., and Sud, C. K. (1999) Detection of genetic alterations in bladder tumors by comparative genomic hybridization and cytogenetic analysis. *Cancer Genet. Cytogenet.* 110, 87-93.
13. Wodicka, L., Dong, H., Mittmann, M., Ho, M. H., and Lockhart, D. J. (1997) Genome-wide expression monitoring in *Saccharomyces cerevisiae*. *Nat. Biotechnol.* 15, 1359-1367.
14. Christensen, M., Sunde, L., Bokund, L., and Orntoft, T. F. (1999) Comparison of three methods of microsatellite detection. *Scand. J. Clin. Lab. Invest.* 59, 167-177.
15. Celis, J. E., Ostergaard, M., Basso, B., Celis, A., Lauridsen, J. B., Ratz, G. P., Andersen, L., Hein, B., Wolf, H., Orntoft, T. F., and Rasmussen, H. H. (1996) Loss of adipocyte-type fatty acid binding protein and other protein biomarkers is associated with progression of human bladder transitional cell carcinomas. *Cancer Res.* 56, 4782-4790.
16. Celis, J. E., Ratz, G., Basso, B., Lauridsen, J. B., and Celis, A. (1994) In *Cell Biology: A Laboratory Handbook* (Celis, J. E., ed) Vol. 3, pp. 222-230, Academic Press, Orlando, FL.
17. Ohlsson, R., Tycko, B., and Sapiezko, C. (1998) Monoallelic expression: 'there can only be one'. *Trends Genet.* 14, 435-438.
18. Hollander, G. A., Zuklys, S., Morel, C., Mizoguchi, E., Mobisson, K., Simpson, S., Terhorst, C., Wishart, W., Golan, D. E., Shan, A. K., and Burakoff, S. J. (1998) Monoallelic expression of the Interleukin-2 locus. *Science* 279, 2118-2121.
19. Brannan, C. J., and Bartolomei, M. S. (1998) Mechanisms of genomic imprinting. *Curr. Opin. Genet. Dev.* 8, 164-170.
20. Ohlsson, R., Cui, H., He, L., Pfeifer, S., Malmkumpu, H., Jiang, S., Feinberg, A. P., and Hedborg, F. (1999) Mosai allelic insulin-like growth factor 2 expression patterns reveal a link between Wilms' tumorigenesis and epigenetic heterogeneity. *Cancer Res.* 59, 3888-3892.
21. Cui, H., Hedborg, F., He, L., Nordenskjold, A., Sandstedt, B., Pfeifer, Ohlsson, S., and Ohlsson, R. (1997) Inactivation of H19, an imprinted and putative tumor repressor gene, is a preneoplastic event during Wilms' tumorigenesis. *Cancer Res.* 57, 4469-4473.
22. Gafitisk, T., Seldanha, A. J., Styles, C. A., Lander, E. S., and Fink, G. R. (1999) Ploidy regulation of gene expression. *Science* 285, 251-254.
23. Tsao, J., Yatabe, Y., Mark, I. D., Hayan, K., Jones, P. A., and Shibata, D. (2000) Bladder cancer genotype stability during clinical progression. *Genes Chromosomes Cancer* 28, 28-32.
24. Zong, C., Schummer, M., Hood, L., and Morris, D. R. (1999) Messenger RNA translation state: the second dimension of high-throughput expression screening. *Proc. Natl. Acad. Sci. U.S.A.* 96, 10632-10638.
25. Anderson, L., and Selthamer, J. (1997) Comparison of selected mRNA and protein abundances in human liver. *Electrophoresis* 18, 533-537.
26. Ideker, T., Thorsson, V., Ransh, J. A., Christmas, R., Buhler, J., Eng, J. K., Bumgarner, R., Goodell, D. R., Aebersold, R., and Hood, L. (2001) Integrated genomic and proteomic analyses of a systematically perturbed metabolic network. *Science* 292, 928-934.

EXHIBIT 3

214: Urol Res. 2000 Oct;28(5):308-15.

Related Articles, Links

 SpringerLink

Expression of cadherins and catenins in paired tumor and non-neoplastic primary prostate cultures and corresponding prostatectomy specimens.

Wang J, Krill D, Torbenson M, Wang Q, Bisceglia M, Stoner J, Thomas A, DeFlavia P, Dhir R, Becich MJ.

Department of Pathology, University of Pittsburgh Medical Center, PA, USA.


Cadherins are a family of transmembrane proteins that play a crucial role in cell differentiation, cell migration, and intercellular adhesion. Cadherins are associated with catenins through their highly conserved cytoplasmic domain. Down-regulation of E-cadherin protein has been shown in various human cancers. This study examined the expression of cadherins and associated catenins at the mRNA level. Paired tumor and nonneoplastic primary prostate cultures were obtained from surgical specimens. Quantitative multiplex fluorescence reverse transcriptase-polymerase chain reaction (QMF RT-PCR) and quantitative analysis were performed and correlated with immunostain results. Six of seven cases of neoplastic cultures showed moderately-to-markedly decreased levels of E-cadherin and P-cadherin mRNA. Similar losses of alpha-catenin and beta-catenin mRNA were also observed. The results of QMF RT-PCR showed good correlation with the results of immunohistochemical studies based on corresponding formalin-fixed sections. In conclusion, this paper presents a coordinated down-regulation in the expression of E-cadherin and associated catenins at the mRNA and protein level in most of the cases studied. This down-regulation may play an important role in the pathogenesis of prostate cancer.

PMID: 11127708 [PubMed - indexed for MEDLINE]

EXHIBIT 4

93: Int J Cancer. 2003 Oct 10;106(6):848-55.

Related Articles, Links



Vascular endothelial growth factor expression correlates with matrix metalloproteinases MT1-MMP, MMP-2 and MMP-9 in human glioblastomas.

Munaut C, Noel A, Hougrand O, Foidart JM, Boniver J, Deprez M.

Laboratory of Tumour and Development Biology, University of Liege, Liege, Belgium.

Vascular endothelial growth factor (VEGF) is the major endothelial mitogen in central nervous system neoplasms and it is expressed in 64-95% of glioblastomas (GBMs). Tumour cells are the main source of VEGF in GBMs whereas VEGF receptors (VEGFR-1, its soluble form sVEGFR-1, VEGFR-2 and neuropilin-1) are expressed predominantly by endothelial cells. Infiltrating tumour cells and newly-formed capillaries progress through the extracellular matrix by local proteolysis involving matrix metalloproteinases (MMPs). Recent studies have shown that VEGF expression and bioavailability can be modulated by MMPs. We reported previously that the expression of MT1-MMP in human breast cancer cells was associated with an enhanced VEGF expression. We used quantitative RT-PCR, Western blot, gelatin zymography and immunohistochemistry to study the expression of VEGF, VEGFR-1, VEGFR-2, sVEGFR-1, neuropilin-1, MT1-MMP, MMP-2, MMP-9 and TIMP-2 in 20 human GBMs and 5 normal brains. The expression of these MMPs was markedly increased in most GBMs with excellent correlation between mRNA and protein levels; activated forms of MMP-2 and MMP-9 were present in 8/18 and 7/18 of GBMs. A majority of GBMs (17/20) also expressed high levels of VEGF, as previously reported, with strong correlation between VEGF and MT1-MMP gene expression levels, and double immunostaining showed that VEGF and MT1-MMP peptides co-localize in tumour and endothelial cells. Our results suggest that the interplay between metalloproteinases and VEGF previously described in experimental tumours may also be operative in human GBMs. Because of its dual ability to activate MMP-2 and to up-regulate VEGF, MT1-MMP might be of central importance in the growth of GBMs and represent an interesting target for anti-cancer treatments. Copyright 2003 Wiley-Liss, Inc.

PMID: 12918061 [PubMed - indexed for MEDLINE]

BEST AVAILABLE COPY

EXHIBIT 5

89: Leuk Lymphoma. 2003 Aug;44(8):1385-94.

Related Articles, Links

Real-time quantitative RT-PCR of cyclin D1 mRNA in mantle cell lymphoma: comparison with FISH and immunohistochemistry.

Hui P, Howe JG, Crouch J, Nimmakayalu M, Oumsiyeh MB, Tallini G, Flynn SD, Smith BR.

Department of Laboratory Medicine, Yale University School of Medicine, 333 Cedar Street, P.O. Box 208035, New Haven, CT 06520-8035, USA.

Presence of the balanced translocation t(11;14)(q13;q32) and the consequent overexpression of cyclin D1 found in mantle cell lymphoma (MCL) has been shown to be of important diagnostic value. Although many molecular and immunohistochemical approaches have been applied to analyze cyclin D1 status, correlative studies to compare different methods for the diagnosis of MCL are lacking. In this study, we examined 39 archived paraffin specimens from patients diagnosed with a variety of lymphoproliferative diseases including nine cases meeting morphologic and immunophenotypic criteria for MCL by: (1) real-time quantitative RT-PCR to evaluate cyclin D1 mRNA expression; (2) dual fluorescence in situ hybridization (FISH) to evaluate the t(11;14) translocation in interphase nuclei; and (3) tissue array immunohistochemistry to evaluate the cyclin D1 protein level. Among the nine cases of possible MCL, seven cases showed overexpression of cyclin D1 mRNA (cyclin D1 positive MCL) and two cases showed no cyclin D1 mRNA increase (cyclin D1 negative "MCL-like"). In six of seven cyclin D1 positive cases, the t(11;14) translocation was demonstrated by FISH analysis; in one case FISH was unsuccessful. Six of the seven cyclin D1 mRNA overexpressing cases showed increased cyclin D1 protein on tissue array immunohistochemistry; one was technically suboptimal. Among the two cyclin D1 negative MCL-like cases, FISH confirmed the absence of the t(11;14) translocation in both cases. All other lymphoproliferative diseases studied were found to have low or no cyclin D1 mRNA expression and were easily distinguishable from the cyclin D1 overexpressing MCLs by all three techniques. In addition, to confirming the need to assess cyclin D1 status, as well as, morphology and immunophenotyping to establish the diagnosis of MCL, this study demonstrates good correlation and comparability between measure of cyclin D1 mRNA, the 11;14 translocation and cyclin D1 protein.

Publication Types:

- Evaluation Studies

PMID: 12952233 [PubMed - indexed for MEDLINE]

BEST AVAILABLE COPY

EXHIBIT 6

8: Int J Biochem Cell Biol. 2005 Oct;37(10):2196-206. Epub 2004 Dec 7.

Related Articles,
Links

ELSEVIER

Increased expression of proteasome subunits in skeletal muscle of cancer patients with weight loss.

Khal J, Hine AV, Fearon KC, Dejong CH, Tisdale MJ.

Pharmaceutical Sciences Research Institute, Aston University, Birmingham B4 7ET, UK.

Atrophy of skeletal muscle is common in patients with cancer and results in increased morbidity and mortality. In order to design effective therapy the mechanism by which this occurs needs to be elucidated. Most studies suggest that the ubiquitin-proteasome proteolytic pathway is most important in intracellular proteolysis, although there have been no reports on the activity of this pathway in patients with different extents of weight loss. In this report the expression of the ubiquitin-proteasome pathway in rectus abdominis muscle has been determined in cancer patients with weight loss of 0-34% using a competitive reverse transcriptase polymerase chain reaction to measure expression of mRNA for proteasome subunits C2 and C5, while protein expression has been determined by western blotting. Overall, both C2 and C5 gene expression was increased by about three-fold in skeletal muscle of cachectic cancer patients (average weight loss 14.5 \pm 2.5%), compared with that in patients without weight loss, with or without cancer. The level of gene expression was dependent on the amount of weight loss, increasing maximally for both proteasome subunits in patients with weight loss of 12-19%. Further increases in weight loss reduced expression of mRNA for both proteasome subunits, although it was still elevated in comparison with patients with no weight loss. There was no evidence for an increase in expression at weight losses less than 10%. There was a good correlation between expression of proteasome 20S α subunits, detected by western blotting, and C2 and C5 mRNA, showing that increased gene expression resulted in increased protein synthesis. Expression of the ubiquitin conjugating enzyme, E2(14k), with weight loss followed a similar pattern to that of proteasome subunits. These results suggest variations in the expression of key components of the ubiquitin-proteasome pathway with weight loss of cancer patients, and suggest that another mechanism of protein degradation must be operative for patients with weight loss less than 10%.

PMID: 16125116 [PubMed - in process]

BEST AVAILABLE COPY

EXHIBIT 7

269: Am J Pathol. 1999 Sep;155(3):815-22.

Related Articles, Links



Id-1 and Id-2 are overexpressed in pancreatic cancer and in dysplastic lesions in chronic pancreatitis.

Maruyama H, Kleeff J, Wildi S, Friess H, Buchler MW, Israel MA, Korc M.

Division of Endocrinology, Department of Medicine, University of California, Irvine, USA.

Id proteins antagonize basic helix-loop-helix proteins, inhibit differentiation, and enhance cell proliferation. In this study we compared the expression of Id-1, Id-2, and Id-3 in the normal pancreas, in pancreatic cancer, and in chronic pancreatitis (CP). Northern blot analysis demonstrated that all three Id mRNA species were expressed at high levels in pancreatic cancer samples by comparison with normal or CP samples. Pancreatic cancer cell lines frequently coexpressed all three Ids, exhibiting a good correlation between Id mRNA and protein levels, as determined by immunoblotting with highly specific anti-Id antibodies. Immunohistochemistry using these antibodies demonstrated the presence of faint Id-1 and Id-2 immunostaining in pancreatic ductal cells in the normal pancreas, whereas Id-3 immunoreactivity ranged from weak to strong. In the cancer tissues, many of the cancer cells exhibited abundant Id-1, Id-2, and Id-3 immunoreactivity. Scoring on the basis of percentage of positive cells and intensity of immunostaining indicated that Id-1 and Id-2 were increased significantly in the cancer cells by comparison with the respective controls. Mild to moderate Id immunoreactivity was also seen in the ductal cells in the CP-like areas adjacent to these cells and in the ductal cells of small and interlobular ducts in CP. In contrast, in dysplastic and atypical papillary ducts in CP, Id-1 and Id-2 immunoreactivity was as significantly elevated as in the cancer cells. These findings suggest that increased Id expression may be associated with enhanced proliferative potential of pancreatic cancer cells and of proliferating or dysplastic ductal cells in CP.

PMID: 10487839 [PubMed - indexed for MEDLINE]

BEST AVAILABLE COPY

EXHIBIT 8

283: Neurosci Lett. 1999 Apr 23;265(3):191-4.

Related Articles, Links

ELSEVIER
ELSEVIER

Alterations in neuropeptide Y levels and Y1 binding sites in the Flinders Sensitive Line rats, a genetic animal model of depression.

Caberlotto L, Jimenez P, Overstreet DH, Hurd YL, Mathe AA, Fuxe K.

Department of Neuroscience, Karolinska Institute, Stockholm, Sweden.

Previously, we observed specific alterations of neuropeptide Y (NPY) and Y1 receptor mRNA expression in discrete regions of the Flinders Sensitive Line rats (FSL), an animal model of depression. In order to clarify the correlation between mRNA expression and protein content, radioimmunoassay and receptor autoradiography were currently performed. In the FSL rats, NPY-like immunoreactivity (NPY-LI) was decreased in the hippocampal CA region, while Y1 binding sites were increased; NPY-LI was increased in the arcuate nucleus. Fluoxetine treatment elevated NPY-LI in the arcuate and anterior cingulate cortex and increased Y1 binding sites in the medial amygdala and occipital cortex in both strains. No differences were found regarding the Y2 binding sites. The results demonstrate a good correlation between NPY peptide and mRNA expression, and sustain the possible involvement of NPY and Y1 receptors in depression.

PMID: 10327163 [PubMed - indexed for MEDLINE]

BEST AVAILABLE COPY

EXHIBIT 9

271: Biol Reprod. 1999 Sep;61(3):776-84.

Related Articles, Links

FREE full text article at
www.biolroprod.org

Follicle-stimulating hormone receptor and its messenger ribonucleic acid are present in the bovine cervix and can regulate cervical prostanoid synthesis.

Mizrachi D, Shemesh M.

Department of Hormone Research, Kimron Veterinary Institute, Bet Dagan, Israel 50250.

The hypothesis that FSH regulates the bovine cervical prostaglandin E(2) (PGE(2)) synthesis that is known to be associated with cervical relaxation and opening at the time of estrus was investigated. Cervical tissue from pre-estrous/estrous, luteal, and postovulatory cows were examined for 1) the presence of bovine (b) FSH receptor (R) and its corresponding mRNA and 2) the effect of FSH on the PGE(2) regulatory pathway in vitro. The presence of bFSHR mRNA in the cervix (maximal during pre-estrus/estrus) was demonstrated by the expression of a reverse transcription (RT) polymerase chain reaction (PCR) product (384 base pairs) specific for bFSHR mRNA and sequencing. Northern blotting revealed three transcripts (2.5, 3.3, and 3.8 kilobases [kb]) in cervix from pre-estrous/estrous cows. The level of FSHR (75 kDa) was significantly higher ($p < 0.01$) in Western blots of pre-estrous/estrous cervix than in other cervical tissues. There was a good correlation between the 75-kDa protein expression and its corresponding transcript of 2.55 kb throughout the estrous cycle as described by Northern blot analysis as well as RT-PCR. Incubation of FSH (10 ng/ml) with pre-estrous/estrous cervix resulted in a 3-fold increase in the expression of FSHR and a 2-fold increase in both G protein ($\alpha(s)$) and cyclooxygenase II. FSH (5-20 ng/ml) significantly increased ($p < 0.01$) cAMP, inositol phosphate ($p < 0.01$), and PGE(2) ($p < 0.01$) production by pre-estrous/estrous cervix but not by cervix at the other stages. We conclude that bovine cervix at the time of the peripheral plasma FSH peak (pre-estrus/estrus) contains high levels of FSHR and responds to FSH by increasing the PGE(2) production responsible for cervical relaxation at estrus.

PMID: 10456856 [PubMed - indexed for MEDLINE]

BEST AVAILABLE COPY

EXHIBIT 10

232: J Urol. 2000 Sep;164(3 Pt 2):1026-30.

Related Articles, Links



The decompensated detrusor III: impact of bladder outlet obstruction on sarcoplasmic endoplasmic reticulum protein and gene expression.

Stein R, Gong C, Hutcheson JC, Canning DA, Zderic SA.

Division of Urology, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, USA.

PURPOSE: Regulation of calcium ion homeostasis has a significant role in smooth muscle contractility. The sarcoplasmic endoplasmic reticulum, calcium, magnesium, adenosine triphosphatase (SERCA) is a regulatory ion pump that may have a role in the functional outcome after outlet obstruction. We investigate what correlation if any existed between SERCA protein and gene expression, and the contractile properties in the same bladder. **MATERIALS AND METHODS:** Standardized partial bladder outlet obstructions were created in adult New Zealand white rabbits, which were divided into control, sham operated and obstructed groups. Muscle strip studies subcategorized the obstructed group into compensated (force greater than 50% of control) and decompensated (force less than 50% of control). Microsomal membrane and total RNA fractions were prepared from the same bladder tissue. Membrane proteins were used for Western blot analysis using a SERCA specific monoclonal antibody, and total RNA was assessed with Northern blot analysis. **RESULTS:** The relative intensities of signals for the Western and Northern blots demonstrated a strong correlation between protein and gene expression. Furthermore there was a strong association between the loss of SERCA messenger RNA and protein expression and loss of bladder function. **CONCLUSIONS:** Bladder contractility after outlet obstruction is influenced in part by smooth muscle cell ability to maintain calcium homeostasis via SERCA. The loss of SERCA protein expression is mediated by down-regulation in gene expression in the same bladder. These data suggest that smooth muscle ion pump gene expression is in part mechanically (pressure work) regulated.

PMID: 10958733 [PubMed - indexed for MEDLINE]

BEST AVAILABLE COPY

EXHIBIT 11

- 141: Zhonghua Jie He He Hu Xi Za Zhi. 2002 Jun;25(6):337-40. [Related Articles](#), [Links](#)

[The pathogenic role of macrophage migration inhibitory factor in acute respiratory distress syndrome]

[Article in Chinese]

Guo Y, Xie C.

Department of Respiratory Medicine, First Affiliated Hospital of Zhongshan University, Guangzhou 510080 China.

OBJECTIVE To investigate the expression and pathogenic role of macrophage migration inhibitory factor (MIF) in human acute respiratory distress syndrome (ARDS). **METHODS** The serum level of MIF in ARDS patients and normal persons were measured by ELISA method. Peripheral blood mononuclear cell (PBMC) MIF expression was determined by flow-cytometry. The expression of MIF mRNA and protein in the lung tissues were detected by using double immuno histochemistry labeling and in situ hybridization. **RESULTS** The serum level of MIF increased significantly in ARDS patients as compared with normal persons ($P < 0.01$). The percentage of PBMC MIF expression was higher in ARDS patients than in normal controls ($P < 0.01$). In situ hybridization and immunohistochemistry showed undetectable or weak MIF mRNA and protein expression in normal lungs. In contrast, there was marked upregulation of MIF mRNA and protein expression in the ARDS lungs. In ARDS macrophages infiltrated the alveolar space and interstitium, most of which also expressed MIF. Infiltrating macrophages were almost restricted to the areas of severe tissue damage. The MIF expression level showed a strong correlation with the number of infiltrating macrophages. **CONCLUSIONS** The serum level of MIF and PBMC MIF expression increased in ARDS patients with enhanced pulmonary MIF expression and macrophage infiltration, which suggests that MIF plays a pivotal role in the pathogenesis of ARDS.

PMID: 12126556 [PubMed - indexed for MEDLINE]

BEST AVAILABLE COPY

EXHIBIT 12

□ 1: Int J Oncol. 2005 Nov;27(5):1257-63.

Related Articles, Links

Expression of human telomerase reverse transcriptase gene and protein, and of estrogen and progesterone receptors, in breast tumors: Preliminary data from neo-adjuvant chemotherapy.

Kammori M, Izumiyama N, Hashimoto M, Nakamura K, Okano T, Kurabayashi R, Naoki H, Honma N, Ogawa T, Kaminishi M, Takubo K.

Division of Breast and Endocrine Surgery, Department of Surgery, Graduate School of Medicine, The University of Tokyo, Bunkyo-ku, Tokyo 113-8655, Japan. kanmori-dis@umin.ac.jp.

Human telomerase reverse transcriptase (hTERT), the catalytic subunit of telomerase, is very closely associated with telomerase activity. Telomerase has been implicated in cellular immortalization and carcinogenesis. In situ detection of hTERT will aid in determining the localization of telomerase-positive cells. The aim of this study was to detect expression of hTERT mRNA, hTERT protein, estrogen receptor (ER) and progesterone receptor (PR) in paraffin-embedded breast tissue samples and to investigate the relationship between hTERT expression and various clinicopathological parameters in breast tumorigenesis. We used in situ hybridization (ISH) to examine hTERT gene expression, and immunohistochemistry (IHC) to examine expression of hTERT protein, ER and PR, in breast tissues including 64 adenocarcinomas, 2 phyllode tumors and their adjacent normal breast tissues. hTERT gene expression was detected by ISH in 56 (88%) carcinomas, but in neither of the 2 phyllode tumors. hTERT protein expression was detected by IHC in 52 (81%) carcinomas, but in neither of the 2 phyllode tumors. Moreover, ER and PR were expressed in 42 (66%) and 42 (66%) carcinomas, respectively, and in neither of the 2 phyllode tumors. In 4 cases of breast carcinoma that strongly expressed hTERT gene and protein before treatment, neo-adjuvant chemotherapy led to disappearance of gene and protein expression in all cases. There was a strong correlation between detection of hTERT gene expression by ISH and of hTERT protein by ICH in tissue specimens from breast tumors. These results suggest that detection of hTERT protein by ICH can be used to distinguish breast cancers as a potential diagnostic and therapeutic marker.

PMID: 16211220 [PubMed - in process]

BEST AVAILABLE COPY

EXHIBIT 13

266: Mol Cell Biol. 1999 Nov;19(11):7357-68.

Related Articles, Links

FREE full text article at
mcb.asm.org

FREE full text article
in PubMed Central

A sampling of the yeast proteome.

Futcher B, Latter GI, Monardo P, McLaughlin CS, Garrels JI.

Cold Spring Harbor Laboratory, Cold Spring Harbor, New York 11724, USA.
futcher@cshl.org

In this study, we examined yeast proteins by two-dimensional (2D) gel electrophoresis and gathered quantitative information from about 1,400 spots. We found that there is an enormous range of protein abundance and, for identified spots, a good correlation between protein abundance, mRNA abundance, and codon bias. For each molecule of well-translated mRNA, there were about 4,000 molecules of protein. The relative abundance of proteins was measured in glucose and ethanol media. Protein turnover was examined and found to be insignificant for abundant proteins. Some phosphoproteins were identified. The behavior of proteins in differential centrifugation experiments was examined. Such experiments with 2D gels can give a global view of the yeast proteome.

PMID: 10523624 [PubMed - indexed for MEDLINE]

BEST AVAILABLE COPY

EXHIBIT 14

304: J Biol Chem. 1998 Aug 14;273(33):21161-8.

Related Articles, Links

FREE full text article at
www.jbc.org

Overexpression of a DEAD box protein (DDX1) in neuroblastoma and retinoblastoma cell lines.

Godbout R, Packer M, Bie W.

Department of Oncology, Cross Cancer Institute and University of Alberta, 11560 University Ave., Edmonton, Alberta T6G1Z2, Canada.

The DEAD box gene, DDX1, is a putative RNA helicase that is co-amplified with MYCN in a subset of retinoblastoma (RB) and neuroblastoma (NB) tumors and cell lines. Although gene amplification usually involves hundreds to thousands of kilobase pairs of DNA, a number of studies suggest that co-amplified genes are only overexpressed if they provide a selective advantage to the cells in which they are amplified. Here, we further characterize DDX1 by identifying its putative transcription and translation initiation sites. We analyze DDX1 protein levels in MYCN/DDX1-amplified NB and RB cell lines using polyclonal antibodies specific to DDX1 and show that there is a good correlation with DDX1 gene copy number, DDX1 transcript levels, and DDX1 protein levels in all cell lines studied. DDX1 protein is found in both the nucleus and cytoplasm of DDX1-amplified lines but is localized primarily to the nucleus of nonamplified cells. Our results indicate that DDX1 may be involved in either the formation or progression of a subset of NB and RB tumors and suggest that DDX1 normally plays a role in the metabolism of RNAs located in the nucleus of the cell.

PMID: 9694872 [PubMed - indexed for MEDLINE]

EXHIBIT 15

150: Virchows Arch. 2002 May;440(5):461-75. Epub 2002 Mar 23. [Related Articles](#), [Links](#)



Expression of somatostatin receptor types 1-5 in 81 cases of gastrointestinal and pancreatic endocrine tumors. A correlative immunohistochemical and reverse-transcriptase polymerase chain reaction analysis.

Papotti M, Bongiovanni M, Volante M, Allia E, Landolfi S, Helboe L, Schindler M, Cole SL, Bussolati G.

Department of Biomedical Sciences and Oncology, University of Turin, Via Santena 7, 10126 Turin, Italy. mauro.papotti@unito.it

Somatostatin receptors (SSTRs) have been extensively mapped in human tumors by means of autoradiography, reverse-transcriptase polymerase chain reaction (RT-PCR), in situ hybridization (ISH) and immunohistochemistry (IHC). We analyzed the SSTR type 1-5 expression by means of RT-PCR and/or IHC in a series of 81 functioning and non-functioning gastroenteropancreatic (GEP) endocrine tumors and related normal tissues. Moreover, we compared the results with clinical, pathological and hormonal features. Forty-six cases (13 intestinal and 33 pancreatic) were studied for SSTR 1-5 expression using RT-PCR, IHC with antibodies to SSTR types 2, 3, 5 and ISH for SSTR2 mRNA. The vast majority of tumors expressed SSTR types 1, 2, 3 and 5, while SSTR4 was detected in a small minority. Due to the good correlation between RT-PCR and IHC data on SSTR types 2, 3, and 5, thirty-five additional GEP endocrine tumors were studied with IHC alone. Pancreatic insulinomas had an heterogeneous SSTR expression, while 100% of somatostatinomas expressed SSTR5 and 100% gastrinomas and glucagonomas expressed SSTR2. Pre-operative biopsy material showed an overlapping immunoreactivity with that of surgical specimens, suggesting that the SSTR status can be detected in the diagnostic work-up. It is concluded that SSTRs 1-5 are heterogeneously expressed in GEP endocrine tumors and that IHC is a reliable tool to detect SSTR types 2, 3 and 5 in surgical and biopsy specimens.

PMID: 12021920 [PubMed - indexed for MEDLINE]

BEST AVAILABLE COPY

EXHIBIT 16

□ 114: Eur J Cancer. 2003 Mar;39(5):691-7.

Related Articles, Links

ELSEVIER
FOUNDED 1840

Expression of deoxycytidine kinase in leukaemic cells compared with solid tumour cell lines, liver metastases and normal liver.

van der Wilt CL, Kroep JR, Loves WJ, Rots MG, Van Groenigen CJ, Kaspers GJ, Peters GJ.

Department of Medical Oncology, VU University Medical Center, Amsterdam, The Netherlands.

Deoxycytidine kinase (dCK) is required for the phosphorylation of several deoxyribonucleoside analogues that are widely employed as chemotherapeutic agents. Examples include cytosine arabinoside (Ara-C) and 2-chlorodeoxyadenosine (CdA) in the treatment of acute myeloid leukaemia (AML) and gemcitabine to treat solid tumours. In this study, expression of dCK mRNA was measured by a competitive template reverse transcriptase polymerase chain reaction (CT RT-PCR) in seven cell lines of different histological origin, 16 childhood and adult AML samples, 10 human liver samples and 11 human liver metastases of colorectal cancer origin. The enzyme activity and protein expression levels of dCK in the cell lines were closely related to the mRNA expression levels ($r=0.75$, $P=0.026$ and $r=0.86$, $P=0.007$). In AML samples, dCK mRNA expression ranged from 1.16 to 35.25 ($\times 10^{-3}$) dCK/beta-actin. In the cell line panel, the range was 2.97-56.9 ($\times 10^{-3}$) dCK/beta-actin of dCK mRNA expression. The enzyme activity in liver metastases was correlated to dCK mRNA expression ($r=0.497$, $P=0.05$). In the liver samples, these were not correlated. dCK mRNA expression showed only a 36-fold range in liver while a 150-fold range was observed in the liver metastases. In addition, dCK activity and mean mRNA levels were 2.5-fold higher in the metastases than in the liver samples. Since dCK is associated with the sensitivity to deoxynucleoside analogues and because of the good correlation between the different dCK measurements in malignant cells and tumours, the CT-RT PCR assay will be useful in the selection of patients that can be treated with deoxycytidine analogues.

PMID: 12628850 [PubMed - indexed for MEDLINE]

BEST AVAILABLE COPY

EXHIBIT 17

67: Regul Pept. 2004 Feb 15;117(2):127-39.

Related Articles, Links

ELSEVIER
Full Text Article

Galanin in pituitary adenomas.

Grenback E, Bjellerup P, Wallerman E, Lundblad L, Anggard A, Ericson K, Aman K, Landry M, Schmidt WE, Hokfelt T, Hulting AL.

Department of Molecular Medicine, Endocrine and Diabetes Unit, Karolinska Hospital, S-17176 Stockholm, Sweden. Eva.Grenback@ks.se

Tumor galanin content was measured in extracts from human pituitary adenomas using a specific RIA method for monitoring human galanin. Twenty-two out of twenty-four tumors contained galanin with notably high levels in corticotroph adenomas, varying levels in clinically inactive tumors, and low levels in GH secreting adenomas. Tumor galanin and ACTH contents were closely correlated in all tumors. In four young patients with microadenomas and highly active Mb Cushing tumor galanin was inversely related to tumor volume. The molecular form of tumor galanin, studied with reverse-phase HPLC, was homogeneous with the majority of tumor galanin coeluting with standard human galanin. In the tumors analysed with in situ hybridization there was a good correlation between galanin peptide levels and galanin mRNA expression. In some tumors galanin mRNA and POMC levels coexisted, in others they were essentially in different cell populations. Levels of plasma galanin-LI were not related to tumor galanin concentration, and galanin levels were in the same range in sinus petrosus close to the pituitary venous drainage as in peripheral blood. Corticotrophin releasing hormone injections in two patients caused ACTH, but no detectable galanin release into sinus petrosus. Our results demonstrate that corticotroph, but not GH adenomas, express high levels of galanin, in addition to ACTH, and that in some tumors both polypeptides are synthesised in the same cell population. However, galanin levels in plasma were not influenced by the tumor galanin content.

PMID: 14700749 [PubMed - indexed for MEDLINE]

BEST AVAILABLE COPY

EXHIBIT 18

50: Blood. 2004 Nov 1;104(9):2936-9. Epub 2004 Jul 8.

Related Articles, Links

Full text article at
www.bloodjournal.org

BCL2 protein expression parallels its mRNA level in normal and malignant B cells.

Shen Y, Iqbal J, Huang JZ, Zhou G, Chan WC.

Department of Pathology and Microbiology, University of Nebraska Medical Center, Omaha, USA.

The regulation of B-cell lymphoma 2 (BCL2) protein expression in germinal center (GC) B cells has been controversial. Previous reports have indicated posttranscriptional regulation plays a dominant role. However, a number of recent studies contradicted these reports. Using real-time polymerase chain reaction (PCR) and Standardized Reverse Transcriptase-PCR (StaRT-PCR), we measured the level of mRNA expression in GC, mantle zone (MNZ), and marginal zone (MGZ) cells from laser capture microdissection. Both quantitative RT-PCR measurements of microdissected GC cells from tonsils showed that GC cells had low expression of BCL2 transcripts commensurate with the low protein expression level. These results are in agreement with microarray studies on fluorescence-activated cell sorter (FACS)-sorted cells and microdissected GC cells. We also examined BCL2 mRNA and protein expression on a series of 30 cases of diffuse large B-cell lymphoma (DLBCL) and found, in general, a good correlation. The results suggested that BCL2 protein expression is regulated at the transcriptional level in normal B cells and in the neoplastic cells in most B-cell lymphoproliferative disorders.

PMID: 15242877 [PubMed - indexed for MEDLINE]

BEST AVAILABLE COPY

EXHIBIT 19

1: Blood. 2005 Dec 15;106(13):4315-21. Epub 2005 Aug 25.

Related Articles, Links

Full text article at
www.bloodjournal.org

Cyclin D1-negative mantle cell lymphoma: a clinicopathologic study based on gene expression profiling.

Fu K, Weisenburger DD, Greiner TC, Dave S, Wright G, Rosenwald A, Chiorazzi M, Iqbal J, Gesk S, Siebert R, De Jong D, Jaffe ES, Wilson WH, Delabie J, Ott G, Dave BJ, Sanger WG, Smith LM, Rimsza L, Braziel RM, Muller-Hermelink HK, Campo E, Gascoyne RD, Staudt LM, Chan WC; Lymphoma/Leukemia Molecular Profiling Project.

Department of Pathology and Microbiology, University of Nebraska Medical Center, 983135.
Nebraska Medical Center, Omaha, NE 68198-3135, USA. kfu@unmc.edu

Cyclin D1 overexpression is believed to be essential in the pathogenesis of mantle cell lymphoma (MCL). Hence, the existence of cyclin D1-negative MCL has been controversial and difficult to substantiate. Our previous gene expression profiling study identified several cases that lacked cyclin D1 expression, but had a gene expression signature typical of MCL. Herein, we report the clinical, pathologic, and genetic features of 6 cases of cyclin D1-negative MCL. All 6 cases exhibited the characteristic morphologic features and the unique gene expression signature of MCL but lacked the t(11;14)(q13; q32) by fluorescence in situ hybridization (FISH) analysis. The tumor cells also failed to express cyclin D1 protein, but instead expressed either cyclin D2 (2 cases) or cyclin D3 (4 cases). There was good correlation between cyclin D protein expression and the corresponding mRNA expression levels by gene expression analysis. Using interphase FISH, we did not detect chromosomal translocations or amplifications involving CCND2 and CCND3 loci in these cases. Patients with cyclin D1-negative MCL were similar clinically to those with cyclin D1-positive MCL. In conclusion, cases of cyclin D1-negative MCL do exist and are part of the spectrum of MCL. Up-regulation of cyclin D2 or D3 may substitute for cyclin D1 in the pathogenesis of MCL.

PMID: 16123218 [PubMed - indexed for MEDLINE]

BEST AVAILABLE COPY

EXHIBIT 20

3: J Gen Virol. 2005 Oct;86(Pt 10):2769-80.

Related Articles, Links

Full text article of
vir.sgmjournals.org

The alpha(v)beta6 integrin-receptor for Foot-and-mouth disease virus is expressed constitutively on the epithelial cells targeted in cattle.

Monaghan P, Gold S, Simpson J, Zhang Z, Weinreb PH, Violette SM, Alexandersen S, Jackson T.

Institute for Animal Health, Pirbright Laboratory, Ash Road, Pirbright, Surrey GU24 0NF, UK.

Field strains of Foot-and-mouth disease virus (FMDV) use a number of alpha(v)-integrins as receptors to initiate infection on cultured cells, and integrins are believed to be the receptors used to target epithelial cells in animals. In this study, immunofluorescence confocal microscopy and real-time RT-PCR were used to investigate expression of two of the integrin receptors of FMDV, alpha(v)beta6 and alpha(v)beta3, within various epithelia targeted by this virus in cattle. These studies show that alpha(v)beta6 is expressed constitutively on the surfaces of epithelial cells at sites where infectious lesions occur during a natural infection, but not at sites where lesions are not normally formed. Expression of alpha(v)beta6 protein at these sites showed a good correlation with the relative abundance of beta6 mRNA. In contrast, alpha(v)beta3 protein was only detected at low levels on the vasculature and not on the epithelial cells of any of the tissues investigated. Together, these data suggest that in cattle, alpha(v)beta6, rather than alpha(v)beta3, serves as the major receptor that determines the tropism of FMDV for the epithelia normally targeted by this virus.

PMID: 16186231 [PubMed - in process]

BEST AVAILABLE COPY

EXHIBIT 21

Neuroscience. 2005;136(1):147-60. Epub 2005 Sep 21.

Related Articles, Links

ELSEVIER
FULL-TEXT ARTICLE

Modulation of the glutamatergic receptors (AMPA and NMDA) and of glutamate vesicular transporter 2 in the rat facial nucleus after axotomy.

Eleore L, Vassias I, Vidal PP, de Waele C.

LNRS (CNRS-Paris V), ESA 7060, Centre Universitaire des Saints-Peres, 45 rue des Saints-Peres, 75270 Paris Cedex 06, France.

Facial nerve axotomy is a good model for studying neuronal plasticity and regeneration in the peripheral nervous system. We investigated in the rat the effect of axotomy on the different subunits of excitatory glutamatergic AMPA (GLuR1-4), NMDA (NR1, NR2A-D) receptors, post-synaptic density 95, vesicular glutamate transporter 2, beta catenin and cadherin. mRNA levels and/or protein production were analyzed 1, 3, 8, 30 and 60 days after facial nerve axotomy by in situ hybridization and immunohistofluorescence. mRNAs coding for the GLuR2-4, NR1, NR2A, B, D subunits of glutamatergic receptors and for post-synaptic density 95, were less abundant after axotomy. The decrease began as early as 1 or 3 days after axotomy; the mRNAs levels were lowest 8 days post-lesion, and returned to normal or near normal 60 days after the lesion. The NR2C subunit mRNAs were not detected in either lesioned or intact facial nuclei.

Immunohistochemistry using specific antibodies against GLuR2-3 subunits and against NR1 confirmed this down-regulation. There was also a large decrease in vesicular glutamate transporter 2 immunostaining in the axotomized facial nuclei at early stages following facial nerve section. In contrast, no decrease of NR2A subunit and of post-synaptic density 95 could be detected at any time following the lesion. beta Catenin and cadherin immunoreactivity pattern changed around the cell body of facial motoneuron by day 3 after axotomy, and then, tends to recover at day post-lesion 60 days. Therefore, our results suggest a high correlation between restoration of nerve/muscle synaptic contact, synaptic structure and function in facial nuclei. To investigate the mechanisms involved in the change of expression of these proteins following axotomy, the facial nerve was perfused with tetrodotoxin for 8 days. The blockade of action potential significantly decreased GLuR2-3, NR1 and NR2A mRNAs in the ipsilateral facial nuclei. Thus, axotomy-induced changes in mRNA abundance seemed to depend partly on disruption of activity.

PMID: 16182453 [PubMed - in process]

BEST AVAILABLE COPY

EXHIBIT 22

SECOND DECLARATION OF PAUL POLAKIS, Ph.D.

I, Paul Polakis, Ph.D., declare and say as follows:

1. I am currently employed by Genentech, Inc. where my job title is Staff Scientist.
2. Since joining Genentech in 1999, one of my primary responsibilities has been leading Genentech's Tumor Antigen Project, which is a large research project with a primary focus on identifying tumor cell markers that find use as targets for both the diagnosis and treatment of cancer in humans.
3. As I stated in my previous Declaration dated May 7, 2004 (attached as Exhibit A), my laboratory has been employing a variety of techniques, including microarray analysis, to identify genes which are differentially expressed in human tumor tissue relative to normal human tissue. The primary purpose of this research is to identify proteins that are abundantly expressed on certain human tumor tissue(s) and that are either (i) not expressed, or (ii) expressed at detectably lower levels, on normal tissue(s).
4. In the course of our research using microarray analysis, we have identified approximately 200 gene transcripts that are present in human tumor tissue at significantly higher levels than in normal human tissue. To date, we have successfully generated antibodies that bind to 31 of the tumor antigen proteins expressed from these differentially expressed gene transcripts and have used these antibodies to quantitatively determine the level of production of these tumor antigen proteins in both human tumor tissue and normal tissue. We have then quantitatively compared the levels of mRNA and protein in both the tumor and normal tissues analyzed. The results of these analyses are attached herewith as Exhibit B. In Exhibit B, "+" means that the mRNA or protein was detectably overexpressed in the tumor tissue relative to normal tissue and "-" means that no detectable overexpression was observed in the tumor tissue relative to normal tissue.
5. As shown in Exhibit B, of the 31 genes identified as being detectably overexpressed in human tumor tissue as compared to normal human tissue at the mRNA level, 28 of them (i.e., greater than 90%) are also detectably overexpressed in human tumor tissue as compared to normal human tissue at the protein level. As such, in the cases where we have been able to quantitatively measure both (i) mRNA and (ii) protein levels in both (i) tumor tissue and (ii) normal tissue, we have observed that in the vast majority of cases, there is a very strong correlation between increases in mRNA expression and increases in the level of protein encoded by that mRNA.

6. Based upon my own experience accumulated in more than 20 years of research, including the data discussed in paragraphs 4-5 above and my knowledge of the relevant scientific literature, it is my considered scientific opinion that for human genes, an increased level of mRNA in a tumor tissue relative to a normal tissue more often than not correlates to a similar increase in abundance of the encoded protein in the tumor tissue relative to the normal tissue. In fact, it remains a generally accepted working assumption in molecular biology that increased mRNA levels are more often than not predictive of elevated levels of the encoded protein. In fact, an entire industry focusing on the research and development of therapeutic antibodies to treat a variety of human diseases, such as cancer, operates on this working assumption.
7. I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information or belief are believed to be true, and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful statements may jeopardize the validity of the application or any patent issued thereon.

Dated: 3-29-06

By: Paul Polakis

Paul Polakis, Ph.D.

DECLARATION OF PAUL POLAKIS, Ph.D.

I, Paul Polakis, Ph.D., declare and say as follows:

1. I was awarded a Ph.D. by the Department of Biochemistry of the Michigan State University in 1984. My scientific Curriculum Vitae is attached to and forms part of this Declaration (Exhibit A).
2. I am currently employed by Genentech, Inc. where my job title is Staff Scientist. Since joining Genentech in 1999, one of my primary responsibilities has been leading Genentech's Tumor Antigen Project, which is a large research project with a primary focus on identifying tumor cell markers that find use as targets for both the diagnosis and treatment of cancer in humans.
3. As part of the Tumor Antigen Project, my laboratory has been analyzing differential expression of various genes in tumor cells relative to normal cells. The purpose of this research is to identify proteins that are abundantly expressed on certain tumor cells and that are either (i) not expressed, or (ii) expressed at lower levels, on corresponding normal cells. We call such differentially expressed proteins "tumor antigen proteins". When such a tumor antigen protein is identified, one can produce an antibody that recognizes and binds to that protein. Such an antibody finds use in the diagnosis of human cancer and may ultimately serve as an effective therapeutic in the treatment of human cancer.
4. In the course of the research conducted by Genentech's Tumor Antigen Project, we have employed a variety of scientific techniques for detecting and studying differential gene expression in human tumor cells relative to normal cells, at genomic DNA, mRNA and protein levels. An important example of one such technique is the well known and widely used technique of microarray analysis which has proven to be extremely useful for the identification of mRNA molecules that are differentially expressed in one tissue or cell type relative to another. In the course of our research using microarray analysis, we have identified approximately 200 gene transcripts that are present in human tumor cells at significantly higher levels than in corresponding normal human cells. To date, we have generated antibodies that bind to about 30 of the tumor antigen proteins expressed from these differentially expressed gene transcripts and have used these antibodies to quantitatively determine the level of production of these tumor antigen proteins in both human cancer cells and corresponding normal cells. We have then compared the levels of mRNA and protein in both the tumor and normal cells analyzed.
5. From the mRNA and protein expression analyses described in paragraph 4 above, we have observed that there is a strong correlation between changes in the level of mRNA present in any particular cell type and the level of protein

expressed from that mRNA in that cell type. In approximately 80% of our observations we have found that increases in the level of a particular mRNA correlates with changes in the level of protein expressed from that mRNA when human tumor cells are compared with their corresponding normal cells.

6. Based upon my own experience accumulated in more than 20 years of research, including the data discussed in paragraphs 4 and 5 above and my knowledge of the relevant scientific literature, it is my considered scientific opinion that for human genes, an increased level of mRNA in a tumor cell relative to a normal cell typically correlates to a similar increase in abundance of the encoded protein in the tumor cell relative to the normal cell. In fact, it remains a central dogma in molecular biology that increased mRNA levels are predictive of corresponding increased levels of the encoded protein. While there have been published reports of genes for which such a correlation does not exist, it is my opinion that such reports are exceptions to the commonly understood general rule that increased mRNA levels are predictive of corresponding increased levels of the encoded protein.

7. I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information or belief are believed to be true, and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful statements may jeopardize the validity of the application or any patent issued thereon.

Dated: 5/07/04

By: Paul Polakis

Paul Polakis, Ph.D.

SV 2031808 v1

CURRICULUM VITAE

PAUL G. POLAKIS
Staff Scientist
Genentech, Inc
1 DNA Way, MS#40
S. San Francisco, CA 94080

EDUCATION:

Ph.D., Biochemistry, Department of Biochemistry,
Michigan State University (1984)

B.S., Biology. College of Natural Science, Michigan State University (1977)

PROFESSIONAL EXPERIENCE:

2002-present	Staff Scientist, Genentech, Inc S. San Francisco, CA
1999- 2002	Senior Scientist, Genentech, Inc., S. San Francisco, CA
1997 -1999	Research Director Onyx Pharmaceuticals, Richmond, CA
1992- 1996	Senior Scientist, Project Leader, Onyx Pharmaceuticals, Richmond, CA
1991-1992	Senior Scientist, Chiron Corporation, Emeryville, CA.
1989-1991	Scientist, Cetus Corporation, Emeryville CA.
1987-1989	Postdoctoral Research Associate, Genentech, Inc., South San Francisco, CA.
1985-1987	Postdoctoral Research Associate, Department of Medicine, Duke University Medical Center, Durham, NC

BEST AVAILABLE COPY

1984-1985

Assistant Professor, Department of Chemistry,
Oberlin College, Oberlin, Ohio

1980-1984

Graduate Research Assistant, Department of
Biochemistry, Michigan State University
East Lansing, Michigan

PUBLICATIONS:

1. Polakis, P. G. and Wilson, J. E. 1982 Purification of a Highly Bindable Rat Brain Hexokinase by High Performance Liquid Chromatography. **Biochem. Biophys. Res. Commun.** 107, 937-943.
2. Polakis, P.G. and Wilson, J. E. 1984 Proteolytic Dissection of Rat Brain Hexokinase: Determination of the Cleavage Pattern during Limited Digestion with Trypsin. **Arch. Biochem. Biophys.** 234, 341-352.
3. Polakis, P. G. and Wilson, J. E. 1985 An Intact Hydrophobic N-Terminal Sequence is Required for the Binding Rat Brain Hexokinase to Mitochondria. **Arch. Biochem. Biophys.** 236, 328-337.
4. Uhing, R.J., Polakis, P.G. and Snyderman, R. 1987 Isolation of GTP-binding Proteins from Myeloid HL60 Cells. **J. Biol. Chem.** 262, 15575-15579.
5. Polakis, P.G., Uhing, R.J. and Snyderman, R. 1988 The Formylpeptide Chemoattractant Receptor Copurifies with a GTP-binding Protein Containing a Distinct 40 kDa Pertussis Toxin Substrate. **J. Biol. Chem.** 263, 4969-4979.
6. Uhing, R. J., Dillon, S., Polakis, P. G., Truett, A. P. and Snyderman, R. 1988 Chemoattractant Receptors and Signal Transduction Processes in Cellular and Molecular Aspects of Inflammation (Poste, G. and Crooke, S. T. eds.) pp 335-379.
7. Polakis, P.G., Evans, T. and Snyderman 1989 Multiple Chromatographic Forms of the Formylpeptide Chemoattractant Receptor and their Relationship to GTP-binding Proteins. **Biochem. Biophys. Res. Commun.** 161, 276-283.
8. Polakis, P. G., Snyderman, R. and Evans, T. 1989 Characterization of G25K, a GTP-binding Protein Containing a Novel Putative Nucleotide Binding Domain. **Biochem. Biophys. Res. Commun.** 160, 25-32.
9. Polakis, P., Weber, R.F., Nevins, B., Didsbury, J. Evans, T. and Snyderman, R. 1989 Identification of the *ral* and *rac1* Gene Products, Low Molecular Mass GTP-binding Proteins from Human Platelets. **J. Biol. Chem.** 264, 16383-16389.
10. Snyderman, R., Perianin, A., Evans, T., Polakis, P. and Didsbury, J. 1989 G Proteins and Neutrophil Function. In ADP-Ribosylating Toxins and G Proteins: Insights into Signal Transduction. (J. Moss and M. Vaughn, eds.) Amer. Soc. Microbiol. pp. 295-323.

11. Hart, M.J., Polakis, P.G., Evans, T. and Cerrione, R.A. 1990 The Identification and Characterization of an Epidermal Growth Factor-Stimulated Phosphorylation of a Specific Low Molecular Mass GTP-binding Protein in a Reconstituted Phospholipid Vesicle System. *J. Biol. Chem.* 265, 5990-6001.
12. Yatani, A., Okabe, K., Polakis, P., Halenbeck, R., McCormick, F. and Brown, A. M. 1990 ras p21 and GAP Inhibit Coupling of Muscarinic Receptors to Atrial K⁺ Channels. *Cell*. 61, 769-776.
13. Munemitsu, S., Innis, M.A., Clark, R., McCormick, F., Ullrich, A. and Polakis, P.G. 1990 Molecular Cloning and Expression of a G25K cDNA, the Human Homolog of the Yeast Cell Cycle Gene CDC42. *Mol. Cell. Biol.* 10, 5977-5982.
14. Polakis, P.G., Rubinfeld, B., Evans, T. and McCormick, F. 1991 Purification of Plasma Membrane-Associated GTPase Activating Protein (GAP) Specific for rap-1/krev-1 from HL60 Cells. *Proc. Natl. Acad. Sci. USA* 88, 239-243.
15. Moran, M. F., Polakis, P., McCormick, F., Pawson, T. and Ellis, C. 1991 Protein Tyrosine Kinases Regulate the Phosphorylation, Protein Interactions, Subcellular Distribution, and Activity of p21ras GTPase Activating Protein. *Mol. Cell. Biol.* 11, 1804-1812.
16. Rubinfeld, B., Wong, G., Bekesi, E., Wood, A., McCormick, F. and Polakis, P. G. 1991 A Synthetic Peptide Corresponding to a Sequence in the GTPase Activating Protein Inhibits p21^{ras} Stimulation and Promotes Guanine Nucleotide Exchange. *Internatl. J. Peptide and Prot. Res.* 38, 47-53.
17. Rubinfeld, B., Munemitsu, S., Clark, R., Conroy, L., Watt, K., Crosier, W., McCormick, F., and Polakis, P. 1991 Molecular Cloning of a GTPase Activating Protein Specific for the Krev-1 Protein p21^{rap1}. *Cell* 65, 1033-1042.
18. Zhang, K., Papageorge, A., G., Martin, P., Vass, W. C., Olah, Z., Polakis, P., McCormick, F. and Lowy, D. R. 1991 Heterogenous Amino Acids in RAS and Rap1A Specifying Sensitivity to GAP Proteins. *Science* 254, 1630-1634.
19. Martin, G., Yatani, A., Clark, R., Polakis, P., Brown, A. M. and McCormick, F. 1992 GAP Domains Responsible for p21^{ras}-dependent Inhibition of Muscarinic Atrial K⁺ Channel Currents. *Science* 255, 192-194.
20. McCormick, F., Martin, G. A., Clark, R., Bollag, G. and Polakis, P. 1992 Regulation of p21ras by GTPase Activating Proteins. Cold Spring Harbor *Symposia on Quantitative Biology*. Vol. 56, 237-241.
21. Pronk, G. B., Polakis, P., Wong, G., deVries-Smits, A. M., Bos J. L. and McCormick, F. 1992 p60^{v-src} Can Associate with and Phosphorylate the p21^{ras} GTPase Activating Protein. *Oncogene* 7,389-394.
22. Polakis P. and McCormick, F. 1992 Interactions Between p21^{ras} Proteins and Their GTPase Activating Proteins. In Cancer Surveys (Franks, L. M., ed.) 12, 25-42.

23. Wong, G., Muller, O., Clark, R., Conroy, L., Moran, M., **Polakis, P.** and McCormick, F. 1992 Molecular cloning and nucleic acid binding properties of the GAP-associated tyrosine phosphoprotein p62. *Cell* 69, 551-558.
24. **Polakis, P.**, Rubinfeld, B. and McCormick, F. 1992 Phosphorylation of rap1GAP in vivo and by cAMP-dependent Kinase and the Cell Cycle p34^{cdc2} Kinase in vitro. *J. Biol. Chem.* 267, 10780-10785.
25. McCabe, P.C., Haubrauck, H., **Polakis, P.**, McCormick, F., and Innis, M. A. 1992 Functional Interactions Between p21^{rap1A} and Components of the Budding pathway of *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 12, 4084-4092.
26. Rubinfeld, B., Crosier, W.J., Albert, I., Conroy, L., Clark, R., McCormick, F. and **Polakis, P.** 1992 Localization of the rap1GAP Catalytic Domain and Sites of Phosphorylation by Mutational Analysis. *Mol. Cell. Biol.* 12, 4634-4642.
27. Ando, S., Kaibuchi, K., Sasaki, K., Hiraoka, T., Nishiyama, T., Mizuno, T., Asada, M., Nunoi, H., Matsuda, I., Matsuura, Y., **Polakis, P.**, McCormick, F. and Takai, Y. 1992 Post-translational processing of rac p21s is important both for their interaction with the GDP/GTP exchange proteins and for their activation of NADPH oxidase. *J. Biol. Chem.* 267, 25709-25713.
28. Janoueix-Lerosey, I., **Polakis, P.**, Tavitian, A. and deGunzberg, J. 1992 Regulation of the GTPase activity of the ras-related rap2 protein. *Biochem. Biophys. Res. Commun.* 189, 455-464.
29. **Polakis, P.** 1993 GAPs Specific for the rap1/Krev-1 Protein. in GTP-binding Proteins: the ras-superfamily. (J.C. LaCale and F. McCormick, eds.) 445-452.
30. **Polakis, P.** and McCormick, F. 1993 Structural requirements for the interaction of p21^{ras} with GAP, exchange factors, and its biological effector target. *J. Biol Chem.* 268, 9157-9160.
31. Rubinfeld, B., Souza, B., Albert, I., Muller, O., Chamberlain, S., Masiarz, F., Munemitsu, S. and **Polakis, P.** 1993 Association of the APC gene product with beta- catenin. *Science* 262, 1731-1734.
32. Weiss, J., Rubinfeld, B., **Polakis, P.**, McCormick, F. Cavenee, W. A. and Arden, K. 1993 The gene for human rap1-GTPase activating protein (rap1GAP) maps to chromosome 1p35-1p36.1. *Cytogenet. Cell Genet.* 66, 18-21.
33. Sato, K. Y., **Polakis, P.**, Haubruck, H., Fasching, C. L., McCormick, F. and Stanbridge, E. J. 1994 Analysis of the tumor suppressor activity of the K-rev gene in human tumor cell lines. *Cancer Res.* 54, 552-559.
34. Janoueix-Lerosey, I., Fontenay, M., Tobelem, G., Tavitian, A., **Polakis, P.** and DeGunzburg, J. 1994 Phosphorylation of rap1GAP during the cell cycle. *Biochem. Biophys. Res. Commun.* 202, 967-975
35. Munemitsu, S., Souza, B., Mueller, O., Albert, I., Rubinfeld, B., and **Polakis, P.** 1994 The APC gene product associates with microtubules in vivo and affects their assembly in vitro. *Cancer Res.* 54, 3676-3681.

36. Rubinfeld, B. and Polakis, P. 1995 Purification of baculovirus produced rap1GAP. **Methods Enz.** 255,31
37. Polakis, P. 1995 Mutations in the APC gene and their implications for protein structure and function. **Current Opinions in Genetics and Development** 5, 66-71
38. Rubinfeld, B., Souza, B., Albert, I., Munemitsu, S. and Polakis P. 1995 The APC protein and E-cadherin form similar but independent complexes with α -catenin, β -catenin and Plakoglobin. **J. Biol. Chem.** 270, 5549-5555
39. Munemitsu, S., Albert, I., Souza, B., Rubinfeld, B., and Polakis, P. 1995 Regulation of intracellular β -catenin levels by the APC tumor suppressor gene. **Proc. Natl. Acad. Sci.** 92, 3046-3050.
40. Lock, P., Fumagalli, S., Polakis, P. McCormick, F. and Courtneidge, S. A. 1996 The human p62 cDNA encodes Sam68 and not the rasGAP-associated p62 protein. **Cell** 84, 23-24.
41. Papkoff, J., Rubinfeld, B., Schryver, B. and Polakis, P. 1996 Wnt-1 regulates free pools of catenins and stabilizes APC-catenin complexes. **Mol. Cell. Biol.** 16, 2128-2134.
42. Rubinfeld, B., Albert, I., Porfiri, E., Fiol, C., Munemitsu, S. and Polakis, P. 1996 Binding of GSK3 β to the APC- β -catenin complex and regulation of complex assembly. **Science** 272, 1023-1026.
43. Munemitsu, S., Albert, I., Rubinfeld, B. and Polakis, P. 1996 Deletion of amino-terminal structure stabilizes β -catenin in vivo and promotes the hyperphosphorylation of the APC tumor suppressor protein. **Mol. Cell. Biol.** 16, 4088-4094.
44. Hart, M. J., Callow, M. G., Sousa, B. and Polakis P. 1996 IQGAP1, a calmodulin binding protein with a rasGAP related domain, is a potential effector for cdc42Hs. **EMBO J.** 15, 2997-3005.
45. Nathke, I. S., Adams, C. L., Polakis, P., Sellin, J. and Nelson, W. J. 1996 The adenomatous polyposis coli (APC) tumor suppressor protein is localized to plasma membrane sites involved in active epithelial cell migration. **J. Cell. Biol.** 134, 165-180.
46. Hart, M. J., Sharma, S., elMasry, N., Qui, R-G., McCabe, P., Polakis, P. and Bollag, G. 1996 Identification of a novel guanine nucleotide exchange factor for the rho GTPase. **J. Biol. Chem.** 271, 25452.
47. Thomas JE, Smith M, Rubinfeld B, Gutowski M, Beckmann RP, and Polakis P. 1996 Subcellular localization and analysis of apparent 180-kDa and 220-kDa proteins of the breast cancer susceptibility gene, BRCA1. **J. Biol. Chem.** 1996 271, 28630-28635
48. Hayashi, S., Rubinfeld, B., Souza, B., Polakis, P., Wieschaus, E., and Levine, A. 1997 A Drosophila homolog of the tumor suppressor adenomatous polyposis coli

down-regulates β -catenin but its zygotic expression is not essential for the regulation of armadillo. **Proc. Natl. Acad. Sci.** 94, 242-247.

49. Vleminckx, K., Rubinfeld, B., Polakis, P. and Gumbiner, B. 1997 The APC tumor suppressor protein induces a new axis in *Xenopus* embryos. **J. Cell. Biol.** 136, 411-420.

50. Rubinfeld, B., Robbins, P., El-Gamil, M., Albert, I., Porfiri, P. and Polakis, P. 1997 Stabilization of β -catenin by genetic defects in melanoma cell lines. **Science** 275, 1790-1792.

51. Polakis, P. The adenomatous polyposis coli (APC) tumor suppressor. 1997 **Biochem. Biophys. Acta**, 1332, F127-F147.

52. Rubinfeld, B., Albert, I., Porfiri, E., Munemitsu, S., and Polakis, P. 1997 Loss of β -catenin regulation by the APC tumor suppressor protein correlates with loss of structure due to common somatic mutations of the gene. **Cancer Res.** 57, 4624-4630.

53. Porfiri, E., Rubinfeld, B., Albert, I., Hovanes, K., Waterman, M., and Polakis, P. 1997 Induction of a β -catenin-LEF-1 complex by wnt-1 and transforming mutants of β -catenin. **Oncogene** 15, 2833-2839.

54. Thomas JE, Smith M, Tonkinson JL, Rubinfeld B, and Polakis P., 1997 Induction of phosphorylation on BRCA1 during the cell cycle and after DNA damage. **Cell Growth Differ.** 8, 801-809.

55. Hart, M., de los Santos, R., Albert, I., Rubinfeld, B., and Polakis P., 1998 Down regulation of β -catenin by human Axin and its association with the adenomatous polyposis coli (APC) tumor suppressor, β -catenin and glycogen synthase kinase 3 β . **Current Biology** 8, 573-581.

56. Polakis, P. 1998 The oncogenic activation of β -catenin. **Current Opinions in Genetics and Development** 9, 15-21

57. Matt Hart, Jean-Paul Concordet, Irina Lassot, Iris Albert, Rico del los Santos, Herve Durand, Christine Perret, Bonnee Rubinfeld, Florence Margottin, Richard Benarous and Paul Polakis. 1999 The F-box protein β -TrCP associates with phosphorylated β -catenin and regulates its activity in the cell. **Current Biology** 9, 207-10.

58. Howard C. Crawford, Barbara M. Fingleton, Bonnee Rubinfeld, Paul Polakis and Lynn M. Matrisian 1999 The metalloproteinase matrilysin is a target of β -catenin transactivation in intestinal tumours. **Oncogene** 18, 2883-91.

59. Meng J, Glick JL, Polakis P, Casey PJ. 1999 Functional interaction between Galpha(z) and Rap1GAP suggests a novel form of cellular cross-talk. **J Biol Chem.** 17, 36663-9

60. Vijayasurian Easwaran, Virginia Song, **Paul Polakis** and Steve Byers 1999 The ubiquitin-proteosome pathway and serine kinase activity modulate APC mediated regulation of β -catenin-LEF signaling. **J. Biol. Chem.** 274(23):16641-5.
- 61 **Polakis P**, Hart M and Rubinfeld B. 1999 Defects in the regulation of beta-catenin in colorectal cancer. **Adv Exp Med Biol.** 470, 23-32
- 62 Shen Z, Batzer A, Koehler JA, **Polakis P**, Schlessinger J, Lydon NB, Moran MF. 1999 Evidence for SH3 domain directed binding and phosphorylation of Sam68 by Src. **Oncogene.** 18, 4647-53
64. Thomas GM, Frame S, Goedert M, Nathke I, **Polakis P**, Cohen P. 1999 A GSK3- binding peptide from FRAT1 selectively inhibits the GSK3-catalysed phosphorylation of axin and beta-catenin. **FEBS Lett.** 458, 247-51.
65. Peifer M, **Polakis P**. 2000 Wnt signaling in oncogenesis and embryogenesis--a look outside the nucleus. **Science** 287,1606-9.
66. **Polakis P**. 2000 Wnt signaling and cancer. **Genes Dev**;14, 1837-1851.
67. Spink KE, **Polakis P**, Weis WI 2000 Structural basis of the Axin-adenomatous polyposis coli interaction. **EMBO J** 19, 2270-2279.
68. Szeto, W., Jiang, W., Tice, D.A., Rubinfeld, B., Hollingshead, P.G., Fong, S.E., Dugger, D.L., Pham, T., Yansura, D.E., Wong, T.A., Grimaldi, J.C., Corpuz, R.T., Singh J.S., Frantz, G.D., Devaux, B., Crowley, C.W., Schwall, R.H., Eberhard, D.A., Rastelli, L., **Polakis, P.** and Pennica, D. 2001 Overexpression of the Retinoic Acid-Responsive Gene Stra6 in Human Cancers and its Synergistic Induction by Wnt-1 and Retinoic Acid. **Cancer Res** 61, 4197-4204.
69. Rubinfeld B, Tice DA, **Polakis P**. 2001 Axin dependent phosphorylation of the adenomatous polyposis coli protein mediated by casein kinase 1 epsilon. **J Biol Chem** 276, 39037-39045.
70. **Polakis P**. 2001 More than one way to skin a catenin. **Cell** 2001 105, 563-566.
71. Tice DA, Soloviev I, **Polakis P**. 2002 Activation of the Wnt Pathway Interferes with Serum Response Element-driven Transcription of Immediate Early Genes. **J Biol. Chem.** 277, 6118-6123.
72. Tice DA, Szeto W, Soloviev I, Rubinfeld B, Fong SE, Dugger DL, Winer J,

Williams PM, Wieand D, Smith V, Schwall RH, Pennica D, **Polakis P**. 2002 Synergistic activation of tumor antigens by wnt-1 signaling and retinoic acid revealed by gene expression profiling. **J Biol Chem**. 277,14329-14335.

73. **Polakis, P**. 2002 Casein kinase I: A wnt'er of disconnect. **Curr. Biol**. 12, R499.

74. Mao, W., Luis, E., Ross, S., Silva, J., Tan, C., Crowley, C., Chui, C., Franz, G., Senter, P., Koeppen, H., **Polakis, P**. 2004 EphB2 as a therapeutic antibody drug target for the treatment of colorectal cancer. **Cancer Res**. 64, 781-788.

75. Shibamoto, S., Winer, J., Williams, M., **Polakis, P**. 2003 A Blockade in Wnt signaling is activated following the differentiation of F9 teratocarcinoma cells. **Exp. Cell Res**. 29211-20.

76. Zhang Y, Eberhard DA, Frantz GD, Dowd P, Wu TD, Zhou Y, Watanabe C, Luoh SM, **Polakis P**, Hillan KJ, Wood WI, Zhang Z. 2004 GEPIS—quantitative gene expression profiling in normal and cancer tissues. **Bioinformatics**, April 8

EXHIBIT B

	tumor mRNA	tumor IHC
UNQ2525	+	+
UNQ2378	+	+
UNQ972	+	-
UNQ97671	+	+
UNQ2964	+	+
UNQ323	+	+
UNQ1655	+	+
UNQ2333	+	+
UNQ9638	+	+
UNQ8209	+	+
UNQ6507	+	+
UNQ8196	+	+
UNQ9109	+	+
UNQ100	+	+
UNQ178	+	+
UNQ1477	+	+
UNQ1839	+	+
UNQ2079	+	+
UNQ8782	+	+
UNQ9646	+	-
UNQ111	+	+
UNQ3079	+	+
UNQ8175	+	+
UNQ9509	+	+
UNQ10978	+	-
UNQ2103	+	+
UNQ1563	+	+
UNQ16188	+	+
UNQ13589	+	+
UNQ1078	+	+
UNQ879	+	+